arXiv:1209.1962v1 [gr-qc] 10 Sep 2012

Averaging and exact perturbations in LTB dust models.

Roberto A. Sussman

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México
(ICN-UNAM), A. P. 70–543, 04510 México D. F., México.

E-mail: ‡`sussman@nucleares.unam.mx`

Abstract. We introduce a scalar weighed average ("q–average") acting on concentric comoving domains in spherically symmetric Lemaître–Tolman–Bondi (LTB) dust models. The resulting averaging formalism allows for an elegant coordinate independent dynamical study of the models, providing as well a valuable theoretical insight on the properties of scalar averaging in inhomogeneous spacetimes. The q–averages of those covariant scalars common to FLRW models (the "q–scalars") identically satisfy FLRW evolution laws and determine for every domain a unique FLRW background state. All curvature and kinematic proper tensors and their invariant contractions are expressible in terms of the q–scalars and their linear and quadratic "local" fluctuations, which convey the effects of inhomogeneity through the ratio of Weyl to Ricci curvature invariants and the magnitude of radial gradients. We define also non–local fluctuations associated with the intuitive notion of a "contrast" with respect to FLRW reference averaged values assigned to a whole domain or time slice. The q–averages of local and non–local quadratic fluctuations are directly and exactly related to statistical correlation moments of the density and Hubble expansion scalar. The evolution equations for the q–scalars and suitably defined perturbations completely determine the dynamics of the models without the back–reaction correlation terms of Buchert's average. A rigorous formalism of exact spherical non–linear perturbations can be defined over a formal FLRW background state associated to the q–scalars, recovering the standard results of linear perturbation theory in the appropriate limit. We briefly explore the possible application of this formalism to open theoretical issues, such as the relation between the growth of inhomogeneity and a definition of gravitational entropy.

PACS numbers: 98.80.-k, 04.20.-q, 95.36.+x, 95.35.+d

1. Introduction.

The spherically symmetric Lemaître–Tolman–Bondi (LTB) dust models [1, 2, 3, 4] are a very popular class of exact solutions of Einstein's equations. Since these models allow us to study non–linear effects in self–gravitating systems by means of analytic expressions or mathematically tractable numerical methods, they have been used in the literature in a wide variety of contexts: as models of cosmological inhomogeneities [5, 6, 7, 8, 9, 10, 11, 12, 13] (see [2, 4, 5, 13] for reviews), as preferred test models in the effort to explain observations without resorting to dark energy [5, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23] (see [23] for a review), cosmic censorship [24, 25] and even in quantum gravity [26].

The LTB models are usually described by their standard original metric variables, which are well suited for most of their physical and cosmological applications, as can be appreciated in the abundant literature (see the book reviews [2, 3, 4]). However, different variables may be better suited to examine the theoretical properties of the models in a coordinate independent manner, for example, the covariant 'fluid flow' scalars in the "1+3" formalism [27, 28, 29, 30, 31, 32] whose evolution equations and spacelike constraints are equivalent to the field equations (see [33] for an innovative approach to the models). Another set of alternative dynamical variables are the "quasi–local" scalars (to be denoted as "q–scalars"), defined as weighed proper volume averages ("q–averages") of covariant fluid flow scalars on spherical comoving domains. The q–scalars can be, either functionals defined on arbitrary fixed domains, or functions ("q–functions") when considering the pointwise dependence of the average on the varying boundary of a domain. By comparing q–scalars with the non–averaged scalars we obtain fluctuations and perturbations, which are exact, not approximated, quantities. The fluctuations and perturbations that can be "local" when the comparison is with q–functions in a pointwise manner, or "non–local" if comparing local non–averaged values with the q–average assigned to a whole domain. The q–functions and their corresponding local perturbations have been applied successfully to examine various aspects and properties of LTB models: to construct an initial value formulation [34], to examine inhomogeneous dark energy sources (quintessence and the Chaplygin gas) [35, 36], to apply a dynamical systems approach [37, 38], to examine their radial asymptotics [39] and the evolution of radial profiles of covariant scalars and void formation [40].

The relevance of applying a formalism of q–averages for the study of LTB models can be appreciated from the fact that averaging has become recently an important open topic in current research in General Relativity. Evidently, as a frame dependent average acting on scalars, the q–average should be examined in reference and comparison to the similar averaging formalism of Buchert [41], which is widely used in the literature (see the comprehensive reviews in [42, 43] and [44, 45] for an alternative covariant averaging formalism acting on proper tensors, see also [46] for further discussion on averaging). Since Buchert's evolution equations reveal the presence of "back–reaction" correlation terms that convey a significant modification of the dynamics, this formalism has a good potential for applications in Cosmology, for example, to understand the effects of non–linearity in structure formation [47], in the re–interpretation of observations [48], in the possibility of explaining cosmic acceleration and dark energy [49], and even in modeling dark energy sources [50]. As physically well motivated toy models of cosmological inhomogeneities, LTB models provide an ideal scenario to apply and test Buchert's formalism [51, 52, 53, 54, 55, 56]

(Buchert's average for LTB models is simply the q–average with unit weight factor). In particular, q–functions and their local fluctuations and perturbations have already been employed as useful tools to examine the existence of back–reaction and "effective" acceleration in generic LTB models in the context of Buchert's formalism [54, 55, 56].

In the present article we address various unresolved issues and extend and enhance previous work. In particular, besides the q–functions and local fluctuations and perturbations that were used in [35, 36, 39, 40, 54, 55, 56], we now consider the full range of q–scalars, which includes (besides q–functions) the q–scalars defined as strict average functionals and their associated non–local fluctuations and perturbations. While some of the background material in sections 3–5 has appeared before, we introduce new elements and further discussion. In particular, we provide in section 4 a more thorough discussion of the properties of all q–scalars, emphasizing their role as LTB scalars that behave as "effective" FLRW scalars, as they (i) satisfy FLRW time dynamics, (ii) mimic FLRW expressions that are widely used in the literature (see void models [14, 15, 16, 17, 18, 19, 20, 21, 22, 23]), and (iii) identify for each domain a unique FLRW background state through Darmois matching conditions (though an actual matching with a FLRW region is optional, not mandatory). We define local and non–local fluctuations and perturbations in section 5 and examine in section 6 their relation with all tensorial objects characteristic of the models (summarized in Appendix D). We show that all proper curvature and kinematic tensors and their invariant contractions are expressible in terms of q–scalars and their local linear and quadratic fluctuations (or perturbations). In particular, the curvature tensors (Riemann, Ricci, Weyl and electric Weyl) and their contractions are entirely expressible in terms of the Ricci scalar, its q–average and and its local fluctuations. While local and non–local quadratic fluctuations of the density and Hubble scalar are different objects, their q–average is the same for every domain (see proof in Appendix C), and as a consequence, the q–average of these fluctuations are averages of invariant scalar contractions and are equal to the statistical variance and covariance (correlation) moments of the density and of the Hubble expansion scalar.

Considering that local perturbations convey the deviation from FLRW geometry through the ratios of Weyl to Ricci curvature invariants and anisotropic (shear) to isotropic expansion, and bearing in mind that these perturbations and their associated q–functions completely describe all proper tensors and scalar contractions, it is natural to expect that these q–scalars and perturbations should also yield a complete and self–consistent system of evolution equations that fully determine the dynamics of the models [35, 36, 37, 38, 39, 40]. In section 7 we derive a similar set of complete evolution equations based on q–averages and non–local perturbations.

In section 8 we show that the back–reaction terms that appear in Buchert's dynamical equations vanish identically in the evolution equations associated with the q–scalars. Hence, the utilization of the q–average (at least as far as LTB models are concerned) leads to different implications from those of Buchert's average: instead of modifying the dynamics by the extra back–reaction terms, the q–scalars provide a more elegant description, with deeper theoretical insight, of the existing fluid flow dynamics, as witnessed by the results of section 6: the direct (non–perturbative) relation between proper tensors and their contractions with local fluctuations and between the q–averages of quadratic fluctuations and statistical correlation moments of the density and Hubble expansion. These simple straightforward theoretically significant relations do not occur with Buchert's averaging (at least in its non–perturbative application to LTB models).

We showed in [35, 36] that a rigorous formalism of exact spherical perturbations on a FLRW background emerges from the q–functions (which define a formal FLRW background) and their local perturbations. We extend this result in section 9 to non–local perturbations, and in sections 10 and 11 by providing a thorough comparison between local and non–local perturbations. In particular, we discuss the cases when local and non–local fluctuations and perturbations are "confined" (*i.e.* localized in a given bounded comoving domain, with or without a matching with a FLRW region), and when they are "asymptotic" (the domain becomes the whole time slice). We also emphasize the fact that local perturbations describe the deviation from FLRW conditions through the local magnitude of radial gradients of covariant scalars, whereas non–local perturbations describe this deviation through the familiar and intuitive notion of a "contrast" between local values of non–averaged scalars and reference average values assigned to a whole domain (or to a whole slice in the asymptotic limit). As a consequence, non–local perturbations are more intuitive, as the sign of their amplitudes corresponds to the familiar positive/negative sign that we associate to over/under densities. This sign is the opposite for local perturbations: over/under densities are negative/positive. In section 12 we show that in the linear limit both perturbations (local and non–local) yield the familiar density perturbation equation of linear theory in the comoving gauge.

As a quick application of the q–average formalism, we consider in section 13 the q–averages in the context of the entropy Leibler–Kullback functional from Information Theory proposed by Buchert and coworkers [42, 57, 58, 59]. We obtain the same conjecture whereby a positive entropy production from this functional follows from a negative statistical correlation between the fluctuations of density and the Hubble expansion scalar (which is the average of an invariant quadratic contraction of the electric Weyl and shear tensors). We show by qualitative arguments (as a preliminary result) that this conjecture is fulfilled in the late time evolution of hyperbolic and elliptic models, but not in the early time evolution when the decaying modes are nonzero (non–simultaneous big bang). A summary and final comments are provided in section 14. The paper contains four brief appendices: Appendix A summarizes the standard analytic solutions of the field equations, Appendix B proves that functions of q–scalars are also q–scalars, Appendix C contains the rigorous proof that local and non–local quadratic fluctuations of the density and Hubble scalar have exactly the same q–average for any domain, and Appendix D provides coordinate independent expressions for all curvature and kinematic proper tensors characteristic of the models.

2. The dynamics of LTB models.

The "Lemaître–Tolman–Bondi (LTB) models" are the well known exact solutions of Einstein's field equations characterized by the LTB metric [1, 2, 3]:

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + \frac{R'^2}{1+E}\,\mathrm{d}r^2 + R^2\left(\mathrm{d}\vartheta^2 + \sin^2\vartheta\,\mathrm{d}\varphi^2\right), \tag{1}$$

where $R = R(t,r)$, $R' = \partial R/\partial r$ and $E = E(r)$ (we have set $G = c = 1$ and r has length units). For a normal geodesic comoving 4–velocity $u^a = \delta^a_0$ and a dust source $T^{ab} = \rho\, u^a u^b$ with rest–mass density ρ the field equations reduce to

$$\dot{R}^2 = \frac{2M}{R} + E, \tag{2a}$$

$$2M' = 8\pi\rho R^2 R', \tag{2b}$$

where $\dot R = u^a R_{,a} = \partial R/\partial t$, and $M = M(r)$ is the "effective" mass (the quasi–local Misner–Sharp mass function [36, 60, 61]).

2.1. Covariant objects.

LTB models are characterized by the following covariant objects associated with u^a, T^{ab} and the projection tensor $h_{ab} = g_{ab} + u_a u_b$:

- the rest-mass energy density given by (2*b*): $\rho = u_a u_b T^{ab}$:
- the Hubble expansion scalar:

$$\mathcal{H} \equiv \frac{\theta}{3}, \qquad \text{with:} \quad \theta = \tilde\nabla_a u^a = h_a^b \nabla_b u^a = \frac{2\dot R}{R} + \frac{\dot R'}{R'}, \tag{3}$$

- the Ricci scalar of the hypersurfaces ${}^3\mathcal{T}[t]$ orthogonal to u^a and marked by constant arbitrary t, whose induced metric is $h_{ab} = g_{ij}\delta^i_a\delta^j_b$ with $i, j = r, \vartheta, \varphi$:

$$\mathcal{K} \equiv \frac{\mathcal{R}^{(3)}}{6}, \qquad \text{with:} \quad \mathcal{R}^{(3)} = -\frac{2(ER)'}{R^2 R'}, \tag{4}$$

- the shear tensor ($\sigma_{ab} = \tilde\nabla_{(a}u_{b)} - (\theta/3)h_{ab}$) and the electric Weyl tensor ($E_{ab} = u^c u^d C_{acbd}$):

$$\sigma_{ab} = \Sigma\, \mathbf{e}_{ab}, \qquad \Sigma = \mathbf{e}_{ab}\sigma^{ab} = -\frac{1}{3}\left(\frac{\dot R'}{R'} - \frac{\dot R}{R}\right), \qquad \sigma_{ab}\sigma^{ab} = 6\Sigma^2, \tag{5}$$

$$E_{ab} = \mathcal{E}\, \mathbf{e}_{ab}, \qquad \mathcal{E} = \mathbf{e}_{ab}E^{ab} = -\frac{M}{R^3} + \frac{4\pi}{3}\rho, \qquad E_{ab}E^{ab} = 6\mathcal{E}^2, \tag{6}$$

  where C_{abcd} is the Weyl tensor and the eigenvalues (Σ and $\mathcal{E}$) of σ_{ab} and E_{ab} are given in the eigenframe $\mathbf{e}_{ab} = h_{ab} - 3n_a n_b$, with $n_a = \sqrt{h_{rr}}\delta^r_a$ being a unit 4–vector orthogonal to u^a and to the orbits of SO(3).

It is evident that the dynamics of LTB models becomes completely determined once the following covariant scalars

$$\{\rho,\, \mathcal{H},\, \Sigma,\, \mathcal{E},\, \mathcal{K}\}, \tag{7}$$

have been computed, which can be done through the metric functions or from suitable evolution equations.

2.2. Analytic solutions vs the "fluid flow" covariant approach.

In most work in the literature using LTB models to tackle different problems and applications (see for example [2, 3, 4, 11]) involves the use of the metric functions that follow from the analytic solutions of the field equation (2*a*), which are provided explicitly in Appendix A (see equations (A.1)–(A.3)). These solutions determine the dynamics of the models, and in particular allow us to compute (implicitly) quantities like R' and $\dot R'$, which are needed to compute the covariant scalars (7) from their forms in (2*b*) and (3)–(6).

However, the dynamics of the models can also be fully determined by finding the scalars (7) through suitable evolution equations, such as those that follow from the "1+3" formalism of Ehlers, Ellis, Bruni, Dunsby and van Ellst [27, 28, 29, 32, 30, 31]:

$$\dot{\mathcal{H}} = -\mathcal{H}^2 - \frac{4\pi}{3}\rho - 2\Sigma^2, \tag{8a}$$

$$\begin{aligned}\dot\rho &= -\,3\,\rho\,\mathcal{H}, &(8b)\\ \dot\Sigma &= -\,2\,\mathcal{H}\,\Sigma+\Sigma^2-\mathcal{E}, &(8c)\\ \dot{\mathcal{E}} &= -\,4\pi\,\rho\Sigma-3\,\mathcal{E}\,(\mathcal{H}+\Sigma)\,, &(8d)\end{aligned}$$

together with the spacelike and Hamiltonian constraints

$$(\Sigma+\mathcal{H})'+3\,\Sigma\,\frac{R'}{R}=0,\qquad \left(\frac{4\pi}{3}\,\rho+\mathcal{E}\right)'+3\,\mathcal{E}\,\frac{R'}{R}=0, \tag{9}$$

$$\mathcal{H}^2=\frac{8\pi}{3}\,\rho-\mathcal{K}+\Sigma^2. \tag{10}$$

Solving this system of partial differential equations is equivalent to working out the models through the field equations $(2a)$–$(2b)$ and the solutions (A.1)–(A.3). However, (7) is not the only nor the most convenient 'fluid flow' scalar representation to handle the models.

3. The q–scalars.

As shown in [34, 35, 36, 39, 40], an alternative coordinate independent "fluid flow" representation for the LTB models is provided by the "quasi–local" or q–scalars and their fluctuations and perturbations. We provide below a rigorous definition of these variables. ‡

3.1. The q–averages.

Consider an arbitrary complete § time slice ${}^3\mathcal{T}[t]$ of an LTB model with metric (1) whose proper volume element is

$$\mathrm{d}\mathcal{V}_p=\sqrt{\det(h_{ab})}\,\mathrm{d}^3x=\frac{R^2R'\sin\vartheta}{\mathcal{F}}\,\mathrm{d}r\,\mathrm{d}\vartheta\,\mathrm{d}\varphi, \tag{11}$$

where (from $(2a)$)

$$\mathcal{F}\equiv\sqrt{1+E}=\left[\dot R^2+\left(1-\frac{2M}{R}\right)\right]^{1/2}. \tag{12}$$

Every ${}^3\mathcal{T}[t]$ can be foliated by the class of compact concentric spherical comoving domains diffeomorphic to $\mathcal{D}[r_b]=\chi[r_b]\times\mathbb{S}^2\subset{}^3\mathcal{T}[t]$, where $\mathbb{S}^2$ is the unit 2–sphere and the closed real interval $\chi[r_b]=\{\bar r\,|\,0\le\bar r\le r_b\}$ is a segment of a radial ray with $\bar r=0$ marking the worldline of the symmetry center (fixed point of the rotation group SO(3)). Following [56], we define

---

‡ See [35, 36] for a discussion on the relation between the definition of q–scalars (q–averages and q–functions in (13) and (18)) to the quasi–local or Misner–Sharp mass when $A=\rho$. The notation and units here are different from those of previous articles: q–scalars were denoted as averages "$\langle A\rangle$" in [34] and as A_* in [35, 36], while the symbol m was used for $(4/3)\pi\rho$ and k for $\mathcal{K}$ in [55, 54, 56, 39, 40]. The notation used in (13) expresses the domain dependence of the q–average by the symbol "$[r_b]$" that univocally identifies any domain $\mathcal{D}[r_b]$ by its boundary: the comoving 2–sphere marked by constant arbitrary $r=r_b$. We use the subindex $_q$ to distinguish the q–average from the "standard" proper volume average functional "$\langle A\rangle_p[r_b]$" of Buchert's formalism, whose weight factor $\mathcal{F}$ is a constant. Both averages coincide only for parabolic models (A.1) for which $\mathcal{F}=1$, $E=0$ and self–similar LTB models for which $\mathcal{F}\neq1$, $E=$ const. (see [56]).

§ See [40, 56] for the case of non–complete slices intersecting a curvature singularity.

The q–average. Let $X(\mathcal{D}[r_b])$ denote the set of all smooth integrable LTB scalar functions in a domain $\mathcal{D}[r_b]$ with r_b fixed but arbitrary marking its boundary. The q–average of a scalar $A \in X(\mathcal{D}[r_b])$ is the linear functional $\langle\ \rangle_q[r_b] : X(\mathcal{D}_[r_b]) \to \mathbb{R}$ such that:

$$A \mapsto \langle A\rangle_q[r_b] = \frac{\int_{\mathcal{D}[r_b]} A\,\mathcal{F}\,\mathrm{d}\mathcal{V}_p}{\int_{\mathcal{D}[r_b]} \mathcal{F}\,\mathrm{d}\mathcal{V}_p} = \frac{\int_0^{r_b} A(\bar r)R^2(\bar r)R'(\bar r)\,\mathrm{d}\bar r}{\int_0^{r_b} R^2(\bar r)R'(\bar r)\,\mathrm{d}\bar r}, \tag{13}$$

where the "weight factor" $\mathcal{F}$ is given by (12).

It is straightforward to evaluate in an arbitrary fixed domain $\mathcal{D}[r_b]$ the q–average of those covariant scalars $A = \rho,\ \mathcal{H},\ \mathcal{K}$ that are common with a FLRW cosmology:

$$\frac{4\pi}{3}\langle\rho\rangle_q[r_b] = \frac{4\pi}{3}\frac{\int_0^{r_b}\rho\,R^2\,R'\mathrm{d}\bar r}{\int_0^{r_b} R^2\,R'\mathrm{d}\bar r} = \frac{2M_b}{R_b^3}, \tag{14}$$

$$\langle\mathcal{H}\rangle_q[r_b] = \frac{\int_0^{r_b}\mathcal{H}\,R^2\,R'\mathrm{d}\bar r}{\int_0^{r_b} R^2\,R'\mathrm{d}\bar r} = \frac{\dot R_b}{R_b}, \tag{15}$$

$$\langle\mathcal{K}\rangle_q[r_b] = \frac{\int_0^{r_b}\mathcal{K}\,R^2\,R'\mathrm{d}\bar r}{\int_0^{r_b} R^2\,R'\mathrm{d}\bar r} = -\frac{E_b}{R_b^2} = \frac{1-\mathcal{F}_b^2}{R_b^2}, \tag{16}$$

where $R_b = R(t, r_b)$, $M_b = M(r_b)$, $E_b = E(r_b)$ (subindex $_b$ will denote evaluation at $r = r_b$), and we simply applied the definition (13) to (2b), (3) and (4). The q–averages (14)–(16) transform the field equation (2a) for every $\mathcal{D}[r_0]$ into

$$\langle\mathcal{H}\rangle_q^2[r_b] = \frac{8\pi}{3}\langle\rho\rangle_q[r_b] - \langle\mathcal{K}\rangle_q[r_b], \tag{17}$$

which is formally identical with the Friedman equation for a dust FLRW model given in terms of the q–averages of the equivalent LTB scalars.

It is important to remark that the weight factor (12) is an invariant quantity, since R and M are invariant scalars in spherically symmetric spacetimes [60]. In fact, in the Newtonian limit ($2M/R \ll 1$) this factor reduces to the total binding energy of comoving dust layers, while in the special relativity limit it becomes the "gamma factor" [61].

3.2. The q–functions.

Since $r = r_b$ in (13) and (14)–(16) is arbitrary, the definition of the q–average functional leads in a natural way to local functions with this corresponding rule. We define then:

The q–functions. For every ${}^3\mathcal{T}[t]$ foliated by concentric domains $\mathcal{D}[r]$ with $r \geq 0$, the q–function associated with any $A \in X(\mathcal{D}[r])$ is the real valued function $A_q : {}^3\mathcal{T}[t] \to \mathbb{R}$ such that

$$A_q(r) = \langle A\rangle_q[r], \tag{18}$$

holds for every $r \geq 0$. Strictly speaking, the q–functions are q–averages that depend on a varying domain boundary.

Evidently, (14)–(16) and (18) imply that the q–functions associated with $A = \rho,\ \mathcal{H},\ \mathcal{K}$ are

$$\frac{4\pi}{3}\rho_q = \frac{2M}{R^3}, \qquad \mathcal{H}_q = \frac{\dot R}{R}, \qquad \mathcal{K}_q = \frac{1-\mathcal{F}^2}{R^2}, \tag{19}$$

$$\mathcal{H}_q^2 = \frac{8\pi}{3}\rho_q - \mathcal{K}_q, \tag{20}$$

where all functions above (save those inside the integral sign) are real valued functions that depend on the upper bound of the integration range $0 \leq \bar{r} \leq r$ for arbitrary r (or the domain boundary for arbitrary domains).

3.3. Functionals vs functions.

The difference between the average functional $\langle A\rangle_q[r_b]$ and its associated q–function $A_q(r)$ is subtle but important (see figures 1a and 1b of references [54, 55, 56]). For every fixed arbitrary domain $\mathcal{D}[r_b] \subset {}^3\mathcal{T}[t]$ they are identical at the boundary $r = r_b$, but they differ for all points $r \neq r_b$ because $\langle A\rangle_q[r_b]$ has been assigned by (13) (as a functional) to the whole of $\mathcal{D}[r_b]$ and must be considered as a constant for points $r < r_b$ inside this domain in each time slice ${}^3\mathcal{T}[t]$, while $A_q(r)$ (as a function) varies smoothly along these inner points (and also along outer points $r > r_b$). As a consequence, the $\langle A\rangle_q[r_b]$ and $A_q(r)$ satisfy the same local derivation rules (23)–(24) at the boundary of each generic domain (see next section), but behave differently when integrated along any fixed domain in the range $0 \leq \bar{r} \leq r_b$. Hence:

$$\langle\, \langle A\rangle_q[r_b]\,\rangle_q[r_b] = \langle A\rangle_q[r_b] \qquad \text{but} \qquad \langle\, A_q(r)\,\rangle_q[r_b] \neq A_{qb} = A_q(r_b), \tag{21}$$

holds for every scalar and every domain. The difference between functionals and functions is clearly illustrated in figure 2.

3.4. Notation.

We will adopt henceforth the following conventions in order to simplify the notation:

- The time dependence of functionals and functions will be omitted unless it is needed to avoid ambiguities. The symbol $\bar{r}$ will be used as a dummy variable inside integral signs.
- Unless stated otherwise, radial dependence given in terms of r (*i.e.* "$M(r)$" or "$\rho_q(r)$" or "$\langle\rho\rangle_q[r]$") or altogether omitted (*i.e.* "M" or "ρ_q" or "$\langle\rho\rangle_q$") will be understood as local point–wise dependence like q–functions, whereas radial dependence in terms of r_b will denote functionals that correspond to a fixed arbitrary domain $\mathcal{D}[r_b]$ and functions that equate to them (as in the right hand sides of (14)–(16)).

4. Properties of the q–scalars.

4.1. Time and radial derivatives.

It is straightforward to show that the following commutating rule holds for every scalar A

$$\frac{\partial}{\partial t}\langle A\rangle_q - \left\langle \frac{\partial A}{\partial t}\right\rangle_q = \langle A\rangle\dot{}_q - \langle \dot{A}\rangle_q = 3\langle \mathcal{H}\, A\rangle_q - 3\langle\mathcal{H}\rangle_q\langle A\rangle_q, \tag{22}$$

where we used (15) and expressed (3) as $3\mathcal{H} = \theta = (R^2R')\dot{}/(R^2R')$. The following useful relations are readily obtained:

$$\langle A\rangle_q' = \frac{\partial}{\partial r}\langle A\rangle_q = \frac{3R'}{R}\,[\,A - \langle A\rangle_q\,], \tag{23}$$

$$\langle A\rangle_q = A(r) - \frac{1}{R^3(r)}\int_0^r A'(\bar{r})\,R^3(\bar{r})\,\mathrm{d}\bar{r}, \tag{24}$$

where (24) follows by integrating (13) by parts. The same properties hold identically for the q–functions A_q.

4.2. Functions of q–scalars.

As we prove in Appendix B (see also Appendix C of [62]), all scalars expressible as functions of q–scalars are themselves q–scalars. As an example, consider the Omega factor (analogous to a FLRW Omega factor) defined as a q–function $\Omega_q = U(\rho_q, \mathcal{H}_q)$ or $\Omega_q = U(\rho_q, \mathcal{K}_q)$ by:

$$\Omega_q = \frac{8\pi\rho_q}{3\mathcal{H}_q^2} = \frac{(8\pi/3)\rho_q}{(8\pi/3)\rho_q - \mathcal{K}_q}, \qquad \Omega_q - 1 = \frac{\mathcal{K}_q}{\mathcal{H}_q^2} = \frac{\mathcal{K}_q}{(8\pi/3)\rho_q - \mathcal{K}_q}, \tag{25}$$

whose associated local scalar is given by (B.3) as

$$\text{“}\Omega\text{”} = \Omega_q \left[1 + \frac{\rho - \rho_q}{\rho_q} - \frac{2(\mathcal{H} - \mathcal{H}_q)}{\mathcal{H}_q}\right], \tag{26}$$

where we used the quotation marks to emphasize that $\Omega \neq 8\pi\rho/(3\mathcal{H}^2) = U(\rho, \mathcal{H})$. Notice that (26) can be expressed as $\Omega = \Omega_q(1 + \delta^{(\Omega)})$ with $\delta^{(\Omega)}$ given by (37).

4.3. The q–scalars satisfy FLRW evolution laws and mimic FLRW expressions.

It is straightforward to show that the q–scalars (whether evaluated as q–functions or as functionals in fixed domains $\mathcal{D}[r_b]$) satisfy FLRW evolution laws: ‖

$$\dot{\rho}_q = -3\mathcal{H}_q\rho_q, \qquad \dot{\mathcal{K}}_q = -2\mathcal{H}_q\mathcal{K}_q \tag{27a}$$

$$\dot{\mathcal{H}}_q = -\mathcal{H}_q^2 - \frac{4\pi}{3}\rho_q, \qquad \dot{\Omega}_q = \Omega_q\left(\Omega_q - 1\right), \tag{27b}$$

which are identical to the evolution equations satisfied by the equivalent covariant scalars $\tilde{\rho}$, $\tilde{\mathcal{H}}$, $\tilde{\mathcal{K}}$, $\tilde{\Omega}$ of a FLRW dust model (a tilde will denote henceforth FLRW objects). This fact evidently singles out the q–scalars as LTB scalars that behave as FLRW scalars (in the sense that they comply with FLRW dynamics).

By parametrizing the metric functions in (1) in terms of their fiducial values at an arbitrary time slice $t = t_0$ and making the coordinate choice $R_0 = r$, we can transform (1) into the “FLRW look alike” metric:

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + a^2(t, r)\left[\frac{\Gamma\,\mathrm{d}r^2}{1 - \mathcal{K}_{q0}\,r^2} + r^2\left(\mathrm{d}\vartheta^2 + \sin^2\vartheta\mathrm{d}\varphi^2\right)\right], \tag{28a}$$

$$a \equiv \frac{R}{R_0} = \frac{R}{r}, \qquad \Gamma = \frac{R'/R}{R_0'/R_0} = \frac{rR'}{R} = 1 + \frac{ra'}{a}, \tag{28b}$$

and as a consequence, the q–scalars (19), (20) and (25) take the following “FLRW look alike” forms:

$$\rho_q = \frac{\rho_{q0}}{a^3}, \qquad \mathcal{K}_q = \frac{\mathcal{K}_{q0}}{a^2} \tag{29a}$$

$$\mathcal{H}_q^2 = \mathcal{H}_{q0}\left[\frac{\Omega_{q0}}{a^3} + \frac{1 - \Omega_{q0}}{a^2}\right], \qquad \Omega_q = \frac{\Omega_{q0}}{\Omega_{q0} + (1 - \Omega_{q0})\,a}, \tag{29b}$$

where the subindex $_0$ indicates evaluation at $t = t_0$. The forms (28a)–(28b) and (29a)–(29b) have been used in most recent articles looking at LTB void models [14, 15, 16, 17, 18, 19, 20, 21, 22, 23], where they are loosely introduced as “convenient” ansatzes without any justification other than their “FLRW look alike” forms, thus ignoring the fact that they can be defined rigorously as q–scalars that emerge from the weighed average (13).

‖ For the functionals $\langle A\rangle_q[r_b]$ the derivatives involved are $\langle A\rangle\dot{}_q[r_b]$, which can be evaluated (at $r = r_b$) either directly from (14)–(16), or with the commutation rule (22) and the forms of the local (non–averaged) scalars in (2b), (3), (4) and (26). If using (22) for computing $\langle\mathcal{H}\rangle\dot{}_q[r_b]$ and $\langle\Omega\rangle\dot{}_q[r_b]$ we also need to use the identities (41) and (42) that are proved in Appendix C.

4.4. The q–scalars associate to each domain a unique FLRW "background state".

For every fluid flow FLRW scalar $\tilde{A} = \tilde{\rho},\, \tilde{\mathcal{H}},\, \tilde{\mathcal{K}},\, \tilde{\Omega}$ and its "FLRW equivalent" LTB q–scalar $A_q = \rho_q,\, \mathcal{H}_q,\, \mathcal{K}_q,\, \Omega_q$, the value $A_{qb} = A_q(t, r_b)$ identifies, for each domain $\mathcal{D}[r_b]$ of an LTB model, a specific FLRW dust model that can be smoothly matched (under Darmois matching conditions) at an arbitrary finite comoving radius $r = r_b$ that also marks the boundary of $\mathcal{D}[r_b]$. Consider a dust FLRW universe with metric

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + \tilde{a}^2(t)\,\left[\frac{\mathrm{d}r^2}{1 - k_0\,\ell_0^{-2} r^2} + r^2\left(\mathrm{d}\vartheta^2 + \sin^2\vartheta\,\mathrm{d}\varphi^2\right)\right], \tag{30}$$

where $k_0 = 0 \pm 1$, ℓ_0 is an arbitrary length scale and $\tilde{a}(t)$ is the dimensionless FLRW scale factor. Darmois conditions (necessary and sufficient) for the smoothness of the matching of (30) with an LTB model along a comoving boundary $r = r_b$ are given by [11, 36]

$$\rho_{qb} = \langle\rho\rangle_q[r_b] = \tilde{\rho}(t) = \frac{\tilde{\rho}_0}{\tilde{a}^3}, \qquad \mathcal{K}_{qb} = \langle\mathcal{K}\rangle_q[r_b] = \tilde{\mathcal{K}}(t) = \frac{\tilde{\mathcal{K}}_0}{\tilde{a}^2}, \tag{31a}$$

$$\mathcal{H}_{qb} = \langle\mathcal{H}\rangle_q[r_b] = \tilde{\mathcal{H}}(t) = \frac{\dot{\tilde{a}}}{\tilde{a}}, \qquad \Omega_{qb} = \langle\Omega\rangle_q[r_b] = \tilde{\Omega}(t) = \frac{8\pi\tilde{\rho}}{3\tilde{\mathcal{H}}^2} \tag{31b}$$

$$R_b = \tilde{a}(t)\,r \qquad \text{or} \quad a_b = \tilde{a}, \tag{31c}$$

where the subindex $_0$ denotes evaluation at a fiducial hypersurface $t = t_0$ and $\tilde{a}(t_0) = 1$ holds. We note that conditions (31a) and (31b) require the continuity of the A_q if we would consider a (hypothetical) smooth matching of a domain $\mathcal{D}[r_b]$ of an LTB model with a FLRW region. This feature is strikingly evident if we use the parametrization (28a)–(28b) and (29a)–(29b) with $\tilde{A}_0 = A_{qb0} = A_q(t_0, r_b)$ holds for $A = \rho,\, \mathcal{H},\, \mathcal{K},\, \Omega$. However, from (35) and (57a)–(57b), it is evident that we have $A_b \neq A_{qb}$ and A'_q, $A'_q \neq 0$ in general, and thus the local scalars (7) and the gradients A' and A'_q do not comply with the matching conditions (31a)–(31c).

It is important to remark that the continuity of the A_q under Darmois matching conditions is simply a formal rigorous procedure to identify for every $\mathcal{D}[r_b]$ of an LTB model a particular FLRW dust model that can be defined as a reference "background state". We use the term "state" to emphasize that this identification does not force us to consider an actual matching with a FLRW region, which would yield a "Swiss cheese" configuration in which the background state becomes also an actual background spacetime. Likewise, the A_q also allow us to define a "FLRW equivalent" region to every domain $\mathcal{D}[r_b]$, as they provide through (31a)–(31c) the values of the FLRW scalars $\tilde{A}$ if the whole domain was replaced by an equivalent spherical comoving section of a FLRW spacetime. Therefore, having determined the FLRW background state, the discontinuity of the local scalars and the gradients A' and A'_q is not problematic if we do not wish to construct an actual Swiss cheese model through a smooth matching at the domain's boundary $r = r_b$. In this latter case we can avoid discontinuities by demanding (besides (31a)–(31c)) also the continuity of local scalars at $r = r_b$ through the following extra supplementary condition:

$$A'_{qb} = 0, \quad \Rightarrow \quad A_b = A_{qb} \quad \text{for} \quad A = \rho,\, \mathcal{K},\, \mathcal{H},\, \Omega, \tag{32}$$

which forces A to coincide with A_q at $r = r_b$, and thus explains the appearance (see panels (b) and (d) of figures 1 and 2) of "humps" (if $A' > 0$) or "bags" (if $A' < 0$) in the radial profiles of local scalars (this has been noted in the local density profiles in Swiss cheese models in LTB void models).

5. Fluctuations and perturbations of q–scalars.

5.1. Local fluctuations and perturbations.

If A and $A_q = \langle A\rangle_q$ are both evaluated as real valued functions on the same arbitrary value r that denotes a varying boundary of concentric domains $\mathcal{D}[r]$ for $r \geq 0$, then a local fluctuation can be defined at each r by the simple pointwise comparison:

$$\mathbf{D}(A)(r) = A(r) - A_q(r) = A(r) - \langle A\rangle_q[r]. \tag{33}$$

By normalizing $\mathbf{D}(A)$ with A_q we obtain the useful dimensionless relative local fluctuations that will be called local "perturbations":

$$\delta^{(A)}(r) = \frac{\mathbf{D}(A)(r)}{A_q(r)} = \frac{A(r) - A_q(r)}{A_q(r)}, \tag{34}$$

which comply (from (23) and (24)) with the following useful relation with radial gradients of A_q and A (also valid for the $\langle A\rangle_q$):

$$\delta^{(A)} = \frac{A_q'/A_q}{3R'/R} = \frac{1}{A_q(r)\,R^3(r)}\int_0^r A'(\bar{r})\,R^3(\bar{r})\,\mathrm{d}\bar{r}, \tag{35}$$

that lead, using (23), (24) and (B.3), to the following linear algebraic relations among the $\delta^{(A)}$:

$$2\delta^{(\mathcal{H})} = \Omega_q\,\delta^{(\rho)} + [1-\Omega_q]\,\delta^{(k)}, \tag{36}$$

$$\delta^{(\Omega)} = \delta^{(\rho)} - 2\delta^{(\mathcal{H})} = (1-\Omega_q)\left(\delta^{(\rho)} - \delta^{(\mathcal{K})}\right), \tag{37}$$

where Ω_q is given by (25) and $\delta^{(\Omega)}$ above is consistent with $\Omega = \Omega_q(1+\delta^{(\Omega)})$ in (26).

5.2. Non–local fluctuations and perturbations.

As opposed to local fluctuations that compare A with $A_q = \langle A\rangle_q$ at the same r, we can define for every fixed domain $\mathcal{D}[r_b]$ non–local fluctuations

$$\mathbf{D}_{\mathrm{NL}}(A)(r, r_b) = A(r) - \langle A\rangle_q[r_b], \tag{38}$$

that compare local values $A(r)$ inside the domain with the q–average (functional) of A, which is a non–local quantity assigned to the whole domain (notice that at every ${}^3\mathcal{T}[t]$ the value $\langle A\rangle_q[r_b]$ is effectively a constant for all $r < r_b$ and a function of t for varying ${}^3\mathcal{T}[t]$). Non–local perturbations are readily defined by

$$\delta^{(A)}_{\mathrm{NL}}(r, r_b) = \frac{\mathbf{D}_{\mathrm{NL}}(A)(r, r_b)}{\langle A\rangle_q[r_b]} = \frac{A(r) - \langle A\rangle_q[r_b]}{\langle A\rangle_q[r_b]}, \qquad 0 \leq r < r_b, \quad \langle A\rangle_q[r_b] \neq 0, \tag{39}$$

and, evidently, do not comply with (35) and the properties that follow thereof (notice that $\partial/\partial r[\delta^{(A)}_{\mathrm{NL}}(r, r_b)] = A'/\langle A\rangle_q[r_b]$).

5.3. Statistical fluctuations.

Since evaluating a q–average for a fixed arbitrary $\mathcal{D}[r_b]$ involves integration through the range $0 \leq \bar{r} \leq r_b$, and the functional $\langle A\rangle_q[r_b]$ is effectively constant in this range at every ${}^3\mathcal{T}[t]$, then, following (21), the q–average of local and non–local linear fluctuations is different:

$$\langle \mathbf{D}(A)\rangle_q[r_b] \neq 0 \quad \text{but} \quad \langle \mathbf{D}_{\mathrm{NL}}(A)\rangle_q[r_b] = 0, \tag{40}$$

which means that the $\mathbf{D}_{\mathrm{NL}}(A)$ are statistical fluctuations, but the $\mathbf{D}(A)$ are not.

However, the average of quadratic combinations of local fluctuations of ρ and $\mathcal{H}$ coincides with the average of the same combinations of non–local fluctuations. As we prove in Appendix C (see also Appendix B of [56] for similar results in Buchert's averages), the following results are valid for A, $B = \rho$, $\mathcal{H}$ and for any domain $\mathcal{D}[r_b]$:

$$\langle\, [\mathbf{D}(A)]^2 \,\rangle_q = \langle\, [\mathbf{D}_{\text{NL}}(A)]^2 \,\rangle_q = \langle A^2\rangle_q - \langle A\rangle_q^2 = \mathbf{Var}_q(A), \tag{41}$$

$$\langle\, \mathbf{D}(A)\,\mathbf{D}(B)\,\rangle_q = \langle\, \mathbf{D}_{\text{NL}}(A)\,\mathbf{D}_{\text{NL}}(B)\,\rangle_q = \langle AB\rangle_q - \langle A\rangle_q\,\langle B\rangle_q = \mathbf{Cov}_q(A,B), \tag{42}$$

where we omitted the $[r_b]$ symbol to simplify notation (A, A_q inside inside the $\langle\quad\rangle_q$ brackets depend on $\bar{r} \le r_b$), whereas $\mathbf{Var}_q(A)$ and $\mathbf{Cov}_q(A)$ are the quadratic variance and covariance (correlation) moments associated with $\langle A\rangle_q[r_b]$.

6. Relation of q–scalars and their fluctuations with curvature and kinematic invariants and tensors.

As shown in Appendix D, all proper curvature and kinematic tensors characteristic of LTB models are expressible in terms of algebraic combinations of g_{ab}, u^a and the eigenframe $\mathbf{e}_{ab}$ defined below (5) and (6), with their scalar coefficients given entirely by four scalar invariants: the Ricci scalar ($\mathcal{R}$), the Newman–Penrose conformal invariant ($\Psi_2 = -\mathcal{E}$), the Hubble expansion $\mathcal{H} = \theta/3$ and the eigenvalue of the shear tensor, Σ (see (3) and (5)). As a consequence, the q–scalars ρ_q, $\mathcal{H}_q$ are coordinate independent quantities, as they are directly related to these four scalar invariants:

$$\frac{4\pi}{3}\rho_q = \frac{1}{6}\mathcal{R} - \Psi_2, \qquad \mathcal{H}_q = \mathcal{H} + \Sigma, \tag{43}$$

where we used (2a), (3)–(6), (14)–(17) and/or (19)–(20), and the fact that $\mathcal{R} = 8\pi\rho$. We remark that these relations hold for their associated q–averages $\langle\rho\rangle_q$, $\langle\mathcal{H}\rangle_q$, $\langle\mathcal{K}\rangle_q$ as functions of a varying domain boundary r. The expressions for other q–scalars, $\mathcal{K}_q$ and Ω_q, in terms of $\mathcal{R}$, Ψ_2, Σ, $\mathcal{H}$ can be found by eliminating these q–scalars from the constraints (20) and (25) and substituting (43).

It is easy to show that the eigenvalues of the shear and electric Weyl tensors are local linear fluctuations over the q–averages (or q–functions) of $\mathcal{H}$ and ρ:

$$\Sigma = -\mathbf{D}(\mathcal{H}) = -\,[\,\mathcal{H} - \mathcal{H}_q\,] \qquad \mathcal{E} = -\frac{4\pi}{3}\mathbf{D}(\rho) = -\frac{4\pi}{3}\,[\,\rho - \rho_q\,], \tag{44}$$

where we applied (19) to (3), (5) and (6). Considering that $\Psi_2 = -\mathcal{E}$ and using (D.4), the curvature and kinematic proper tensors (see Appendix D) are expressible in terms of ρ, $\mathcal{H}$ and their fluctuations $\mathbf{D}(\rho)$, $\mathbf{D}(\mathcal{H})$:

$$\mathcal{R}^{ab}_{cd} = \frac{8\pi}{3}\rho\left(3\delta^{[a}_{[c}\delta^{b]}_{d]} + 6\delta^{[a}_{[c}u^{b]}u_{d]} - \delta^{a}_{[c}\delta^{b}_{d]}\right) - \frac{4\pi}{3}\mathbf{D}(\rho)\,\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\,\mathbf{e}^{b]}_{d]}, \tag{45a}$$

$$\mathcal{R}^a_b = 4\pi\rho\,(h^a_b + u^a u_b)\,, \tag{45b}$$

$$E_{ab} = -\frac{4\pi}{3}\mathbf{D}(\rho)\,\mathbf{e}_{ab}, \qquad C^{ab}_{cd} = -\frac{4\pi}{3}\mathbf{D}(\rho)\,\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\,\mathbf{e}^{b]}_{d]}, \tag{45c}$$

$$\sigma_{ab} = -\mathbf{D}(\mathcal{H})\,\mathbf{e}_{ab}, \qquad \mathcal{H}_{ab} = \mathcal{H}h_{ab} - \mathbf{D}(\mathcal{H})\,\mathbf{e}_{ab}, \tag{45d}$$

while their scalar contractions take the form:

$$\mathcal{R}_{abcd}\mathcal{R}^{abcd} = \frac{256\pi^2}{3}\left([\mathbf{D}(\rho)]^2 + \frac{5}{4}\rho^2\right), \qquad \mathcal{R}_{ab}\mathcal{R}^{ab} = 64\pi^2\rho^2, \tag{46a}$$

$$C_{abcd}C^{abcd} = \frac{256\pi^2}{3}[\mathbf{D}(\rho)]^2 = 8E_{ab}E^{ab}, \tag{46b}$$

$$\sigma_{ab}\sigma^{ab} = 6[\mathbf{D}(\mathcal{H})]^2, \qquad \sigma_{ab}E^{ab} = \frac{4\pi}{3}\mathbf{D}(\rho)\mathbf{D}(\mathcal{H}), \tag{46c}$$

which depend exclusively on ρ, $\mathcal{H}$ and their fluctuations. Considering that we have $A = A_q(1+\delta^{(A)})$ and $\mathbf{D}(A) = A_q\delta^{(A)}$ for all A, the tensors (45a)–(45d) and their contractions (46a)–(46c) can be entirely given in terms of ρ_q, $\mathcal{H}_q$ and their local perturbations:

$$E^a_b = -\frac{4\pi}{3}\rho_q\delta^{(\rho)}\,\mathbf{e}^a_b, \qquad \mathcal{R}^a_b = 4\pi\rho_q(1+\delta^{(\rho)})\,(h^a_b + u^a u_b)\,, \tag{47a}$$

$$C^{ab}_{cd} = -\frac{4\pi}{3}\rho_q\delta^{(\rho)}\,\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\,\mathbf{e}^{b]}_{d]}, \tag{47b}$$

$$\mathcal{R}^{ab}_{cd} = \frac{8\pi}{3}\rho_q(1+\delta^{(\rho)})\left(3\delta^{[a}_{[c}\delta^{b]}_{d]} + 6\delta^{[a}_{[c}u^{b]}u_{d]} - \delta^a_{[c}\delta^b_{d]}\right) - \frac{4\pi}{3}\rho_q\delta^{(\rho)}\,\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\,\mathbf{e}^{b]}_{d]}, \tag{47c}$$

$$\sigma^a_b = -\mathcal{H}_q\delta^{(\mathcal{H})}\,\mathbf{e}^a_b, \qquad \mathcal{H}^a_b = \mathcal{H}_q\left[h^a_b + (h^a_b - \mathbf{e}^a_b)\,\delta^{(\mathcal{H})}\right], \tag{47d}$$

$$\mathcal{R}_{abcd}\mathcal{R}^{abcd} = \frac{256\pi^2}{3}\rho_q^2\left([\delta^{(\rho)}]^2 + \frac{5}{4}[1+\delta^{(\rho)}]^2\right), \tag{47e}$$

$$C_{abcd}C^{abcd} = \frac{256\pi^2}{3}[\rho_q\,\delta^{(\rho)}]^2 = 8E_{ab}E^{ab}, \tag{47f}$$

$$\sigma_{ab}\sigma^{ab} = 6[\mathcal{H}_q\,\delta^{(\mathcal{H})}]^2, \qquad \mathcal{H}_{ab}\mathcal{H}^{ab} = \mathcal{H}_q^2\left[1 + (2+3\delta^{(\mathcal{H})})\,\delta^{(\mathcal{H})}\right], \tag{47g}$$

which shows that the variables $\{\rho_q,\ \mathcal{H}_q,\ \delta^{(\rho)},\ \delta^{(\mathcal{H})}\}$ should provide a full dynamical representation for the models, so that $\delta^{(\rho)} = \delta^{(\mathcal{H})} = 0$ indicates a FLRW limit characterized by the vanishing of the shear tensor and Weyl part of the curvature tensors. Notice that all these tensors can also be expressed in terms of other q–scalars ($\mathcal{K}_q$, Ω_q) and their local perturbations ($\delta^{(\mathcal{K})}$, $\delta^{(\Omega)}$), as the latter can be obtained from ρ_q, $\mathcal{H}_q$, $\delta^{(\rho)}$, $\delta^{(\mathcal{H})}$ by means of the algebraic constraints (20), (25), (36) and (37).

Bearing in mind that $\mathcal{R}_q = 8\pi\rho_q$, we can express the conformal invariant Ψ_2 as a local fluctuation of the Ricci scalar: $\Psi_2 = (1/6)\mathbf{D}(\mathcal{R}) = (4\pi/3)\,\mathbf{D}(\rho)$, and thus express curvature tensors (the electric Weyl, Weyl and Riemann) and their contractions in terms of the the Ricci scalar and its fluctuations:

$$E_{ab} = -\frac{1}{6}\mathbf{D}(\mathcal{R})\,\mathbf{e}_{ab}, \qquad C^{ab}_{cd} = -\frac{1}{6}\mathbf{D}(\mathcal{R})\,\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\,\mathbf{e}^{b]}_{d]}, \tag{48a}$$

$$\mathcal{R}^{ab}_{cd} = \frac{1}{3}\mathcal{R}\left(3\delta^{[a}_{[c}\delta^{b]}_{d]} + 6\delta^{[a}_{[c}u^{b]}u_{d]} - \delta^a_{[c}\delta^b_{d]}\right) - \frac{1}{6}\mathbf{D}(\mathcal{R})\,\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\,\mathbf{e}^{b]}_{d]}, \tag{48b}$$

$$\mathcal{R}_{abcd}\mathcal{R}^{abcd} = \frac{4}{3}\left([\mathbf{D}(\mathcal{R})]^2 + \frac{5}{4}\mathcal{R}^2\right), \qquad C_{abcd}C^{abcd} = \frac{4}{3}\,[\mathbf{D}(\mathcal{R})]^2 = 8E_{ab}E^{ab}, \tag{48c}$$

where we used (48a) and (D.3). These expressions provide an elegant geometric interpretation of the inhomogeneity associated with the Weyl part of curvature tensors and scalars in terms of fluctuations of the Ricci scalar.

Since a non–vanishing shear and Weyl curvature are indicative of inhomogeneity, it is very useful to express the basic local perturbations $\delta^{(\rho)}$ and $\delta^{(\mathcal{H})}$ as

$$\delta^{(\rho)} = \frac{\rho - \rho_q}{\rho_q} = -\frac{\mathcal{E}}{4\pi\rho_q/3} = \frac{6\Psi_2}{\mathcal{R} - 6\Psi_2} = \frac{\xi}{1-\xi}, \tag{49a}$$

$$\delta^{(\mathcal{H})} = \frac{\mathcal{H} - \mathcal{H}_q}{\mathcal{H}_q} = -\frac{\Sigma}{\mathcal{H}_q} = -\frac{\Sigma}{\mathcal{H}+\Sigma} = -\frac{\zeta}{1+\zeta}, \tag{49b}$$

where the quotients

$$\xi \equiv \frac{6\Psi_2}{\mathcal{R}}, \qquad \zeta \equiv \frac{\Sigma}{\mathcal{H}}, \tag{50}$$

provide an invariant measure of inhomogeneity through the ratio of Weyl to Ricci scalar curvatures (ξ) and the ratio of anisotropic to isotropic expansion (ζ) ¶ . The remaining perturbations $\delta^{(\mathcal{K})}$ and $\delta^{(\Omega)}$ can be expressed in terms of ξ and ζ by using the constraints (36) and (37) to eliminate them in terms of $\delta^{(\rho)}$ and $\delta^{(\mathcal{H})}$ and then substituting (49*a*) and (49*b*).

Since Σ and $\mathcal{E}$ are local linear fluctuations of $\mathcal{H}$ and ρ, we have for all r_0:

$$\langle\Sigma\rangle_q[r_b] \neq 0, \qquad \langle\mathcal{E}\rangle_q[r_b] \neq 0, \tag{51}$$

However, as a consequence of (41), (42) and (44), the averages of the scalars Σ^2, $\mathcal{E}^2$ and $\Sigma\,\mathcal{E}$ are averages of local quadratic fluctuations, leading to the following statistical fluctuations that relate to the variance moment with respect to $\langle\mathcal{H}\rangle_q$ and $\langle\rho\rangle_q$ at any domain:

$$\langle\Sigma^2\rangle_q = \langle\,[\mathbf{D}(H)]^2\,\rangle_q = \langle\mathcal{H}^2\rangle_q - \langle\mathcal{H}\rangle_q^2 = \mathbf{Var}_q(\mathcal{H}), \tag{52a}$$

$$\langle\mathcal{E}^2\rangle_q = \left[\frac{4\pi}{3}\right]^2\langle\,[\mathbf{D}(\rho)]^2\,\rangle_q = \left[\frac{4\pi}{3}\right]^2\left[\langle\rho^2\rangle_q - \langle\rho\rangle_q^2\right] = \left[\frac{4\pi}{3}\right]^2\mathbf{Var}_q(\rho), \tag{52b}$$

$$\langle\mathcal{E}\,\Sigma\rangle_q = \frac{4\pi}{3}\langle\mathbf{D}(\rho)\,\mathbf{D}(\mathcal{H})\rangle_q = \frac{4\pi}{3}\left[\,\langle\rho\,\mathcal{H}\rangle_q - \langle\rho\rangle_q\,\langle\mathcal{H}\rangle_q\right] = \frac{4\pi}{3}\,\mathbf{Cov}_q(\rho,\mathcal{H}), \tag{52c}$$

so that, in view of (5) and (46*c*), these fluctuations relate to averages of quadratic contractions of the shear and electric Weyl tensors:

$$\langle\sigma_{ab}\sigma^{ab}\rangle_q = 6\langle\Sigma^2\rangle_q = 6\mathbf{Var}_q(\mathcal{H}), \tag{53a}$$

$$\langle E_{ab}E^{ab}\rangle_q = 6\langle\mathcal{E}^2\rangle_q = 6\langle(\Psi_2)^2\rangle_q = \frac{32\pi^2}{3}\mathbf{Var}_q(\rho), \tag{53b}$$

$$\langle\sigma_{ab}E^{ab}\rangle_q = 6\langle\Sigma\,\mathcal{E}\rangle_q = 8\pi\mathbf{Cov}_q(\rho,\mathcal{H}), \tag{53c}$$

while the q–averages of contractions of the Riemann and Weyl tensors are expressible either in terms of ρ or $\mathcal{R}$ and their statistical variance moments:

$$\langle\mathcal{R}_{abcd}\mathcal{R}^{abcd}\rangle_q = \frac{256\pi^2}{3}\left[\mathbf{Var}_q(\rho) + \frac{5}{4}\langle\rho^2\rangle_q\right] = \frac{4}{3}\left[\mathbf{Var}_q(\mathcal{R}) + \frac{5}{4}\langle\mathcal{R}^2\rangle_q\right], \tag{54a}$$

$$\langle C_{abcd}C^{abcd}\rangle_q = \frac{256\pi^2}{3}\,\mathbf{Var}_q(\rho) = \frac{4}{3}\,\mathbf{Var}_q(\mathcal{R}) = 8\langle E_{ab}E^{ab}\rangle_q, \tag{54b}$$

where we used the fact that $\langle\mathcal{R}\rangle_q = 8\pi\langle\rho\rangle_q$ and $\mathcal{R}_{ab}\mathcal{R}^{ab} = \mathcal{R}^2$. We can express the quadratic ratio of Weyl to Ricci curvatures as a sort of "standard deviation" of ρ with respect to $\langle\rho\rangle_q$:

$$\frac{6\langle E_{ab}E^{ab}\rangle_q}{\langle\mathcal{R}_{ab}\mathcal{R}^{ab}\rangle_q} = \frac{6\langle(\Psi_2)^2\rangle_q}{\langle(\mathcal{R})^2\rangle_q} = \frac{6\mathbf{Var}_q(\rho)}{\langle\rho^2\rangle_q} = \frac{\langle\rho^2\rangle_q - \langle\rho\rangle_q^2}{\langle\rho^2\rangle_q}, \tag{55}$$

A similar standard deviation of $\mathcal{H}$ with respect to $\langle\mathcal{H}\rangle_q$ follows as the quotient of averages of quadratic covariant scalars $\sigma_{ab}\sigma^{ab}$ and $\mathcal{H}^2 = \theta^2/9$:

$$\frac{\langle\sigma_{ab}\sigma^{ab}\rangle_q}{6\langle\mathcal{H}^2\rangle_q} = \frac{\mathbf{Var}_q(\mathcal{H})}{\langle\mathcal{H}^2\rangle_q} = \frac{\langle\mathcal{H}^2\rangle_q - \langle\mathcal{H}\rangle_q^2}{\langle\mathcal{H}^2\rangle_q}, \tag{56}$$

where we used (5) and (44).

¶ Wainwight and Andrews (reference [33]) also emphasize the role of the ratio of Weyl to Ricci curvature as an invariant measure of inhomogeneity in LTB models. However, their study was not based on averaging and they consider a quadratic positive definite ratio $C_{abcd}C^{abcd}/\mathcal{R}_{ab}\mathcal{R}^{ab}$. As a contrast, the sign of first order ratio $\Psi_2/\mathcal{R}$ that we use here is not *a priori* defined.

7. Evolution equations.

While the q–scalars A_q or $\langle A\rangle_q[r_b]$ satisfy FLRW evolution laws, such as $(27a)$–$(27b)$, these scalars are not fully determined by these FLRW equations because, unlike their equivalent FLRW scalars $\tilde{A}$, they either depend directly on r or r_b. The missing dynamical information is provided by the evolution equations for the $\delta^{(A)}$ and the $\delta^{(A)}_{\mathrm{NL}}$.

7.1. Local perturbations.

Since all the local covariant scalars (7) (as well the tensors and scalar invariants in $(47a)$–$(47g)$) can be expressed in terms of q–scalars and their local perturbations:

$$\rho = \rho_q\left[1+\delta^{(\rho)}\right], \quad \mathcal{H} = \mathcal{H}_q\left[1+\delta^{(\mathcal{H})}\right], \quad \mathcal{K} = \mathcal{K}_q\left[1+\delta^{(k)}\right], \tag{57a}$$

$$\Sigma = -\mathcal{H}_q\,\delta^{(\mathcal{H})}, \qquad \mathcal{E} = -\frac{4\pi}{3}\rho_q\delta^{(\rho)}, \tag{57b}$$

the evolution equations for the variables ρ_q, $\mathcal{H}_q$, $\delta^{(\rho)}$, $\delta^{(\mathcal{H})}$ should yield a self–consistent and complete set of evolution equations for the LTB models. This system follows readily by inserting $(57a)$ and $(57b)$ into the "1+3" system $(8a)$–$(8d)$ and its constraints (9)–(10). The result is the evolution equations

$$\dot{\rho}_q = -\,3\rho_q\mathcal{H}_q, \tag{58a}$$

$$\dot{\mathcal{H}}_q = -\,\mathcal{H}_q^2 - \frac{4\pi}{3}\rho_q, \tag{58b}$$

$$\dot{\delta}^{(\rho)} = -\,3(1+\delta^{(\rho)})\,\mathcal{H}_q\delta^{(\mathcal{H})}, \tag{58c}$$

$$\dot{\delta}^{(\mathcal{H})} = -\,(1+3\delta^{(\mathcal{H})})\,\mathcal{H}_q\delta^{(\mathcal{H})} + \frac{4\pi\,\rho_q}{3\mathcal{H}_q}(\delta^{(\mathcal{H})}-\delta^{(\rho)}), \tag{58d}$$

plus the algebraic constraints

$$\mathcal{H}_q^2 = \frac{8\pi}{3}\rho_q - \mathcal{K}_q, \quad 2\delta^{(\mathcal{H})} = \Omega_q\,\delta^{(\rho)} + [1-\Omega_q]\,\delta^{(k)}, \quad \delta^{(\Omega)} = \delta^{(\rho)} - 2\delta^{(\mathcal{H})}, \tag{59}$$

that exactly coincide with the general relations (20), (36) and (37), hence they hold at all t (*i.e.* they propagate in time). The following points are worth remarking:

- The constraints (9) of the 1+3 system $(8a)$–$(8d)$ are satisfied trivially from (35) applied to $\delta^{(\rho)}$ and $\delta^{(\mathcal{H})}$.
- The first two constraints in (59) follow from substituting $(57b)$ into (10) and using $(2a)$ and (19), while the third one is obtained by differentiating (25) with respect to r and applying (35). Once $(58a)$–$(58d)$ is solved these constrains and (25) allow us to compute Ω_q, $\mathcal{K}_q$, $\delta^{(k)}$, $\delta^{(\Omega)}$.
- The fact that the constraints (59) of the system $(58a)$–$(58d)$ are algebraic implies a great simplification of the numeric treatment of these fluid flow evolution equations, as they can be effectively integrated as a system of autonomous ODE's in which the initial conditions are restricted by the algebraic constraints, and thus it is far easier to handle than the 1+3 system $(8a)$–$(8d)$ in which the constraints (9) are partial differential equations on r that must be solved before dealing with the time equations.

Considering the relation between the A_q and $\delta^{(A)}$ in the constraints (59), each of the following combination of four scalars:

$$\begin{aligned}&\text{any two of} && A_q = \rho_q,\, \mathcal{H}_q,\, \mathcal{K}_q,\, \Omega_q, && (60a)\\ &\text{plus} && && \\ &\text{any two of} && \delta^{(A)} = \delta^{(\rho)},\, \delta^{(\mathcal{H})},\, \delta^{(\mathcal{K})},\, \delta^{(\Omega)}, && (60b)\end{aligned}$$

provides a full covariant scalar representation for the models, since the remaining pairs of A_q and $\delta^{(A)}$ can be obtained from these algebraic constraints. Each of these representations yields a self–consistent and complete set of evolution equations that is alternative (and equivalent) to that of local covariant scalars given by (7), and thus, completely determine the dynamics of the models.

The evolution equations (58a)–(58d) correspond to the representation $\{\rho_q,\, \mathcal{H}_q,\, \delta^{(\rho)},\, \delta^{(\mathcal{H})}\}$, which is useful to compare with the spherical collapse model and perturbative scenarios of structure formation that consider the density and Hubble velocity as dynamical variables. However, a more appropriate representation for cosmological applications (for example void models) is furnished by the scalars $\{\Omega_q,\, \mathcal{H}_q,\, \delta^{(\Omega)},\, \delta^{(\mathcal{H})}\}$.

7.2. Non–local perturbations.

If we consider non–local perturbations (39) in an arbitrary fixed domain $\mathcal{D}[r_b]$, then the only local scalars in (7) that can be expressed in terms of the variables $\{\langle A\rangle_q[r_b],\, \delta_{\rm NL}^{(A)}\}$ in a similar manner as in (57a) are:

$$\rho = \langle\rho\rangle_q[r_b]\,\left[1+\delta_{\rm NL}^{(\rho)}\right], \quad \mathcal{H} = \langle\mathcal{H}\rangle_q[r_b]\,\left[1+\delta_{\rm NL}^{(\mathcal{H})}\right], \quad \mathcal{K} = \langle\mathcal{K}\rangle_q[r_b]\,\left[1+\delta_{\rm NL}^{(\mathcal{K})}\right]. \tag{61}$$

The remaining local scalars, Σ and $\mathcal{E}$, given by local fluctuations by (44), cannot be expressed as (57b):

$$\Sigma(r) = \mathcal{H}_q(r) - \langle\mathcal{H}\rangle_q[r_b](1+\delta_{\rm NL}^{(\mathcal{H})}), \qquad \mathcal{E}(r) = \frac{4\pi}{3}\left[\rho_q(r) - \langle\rho\rangle_q[r_b]\delta_{\rm NL}^{(\rho)}\right], \tag{62}$$

and as a consequence, the variables $\{\langle A\rangle_q[r_b],\, \delta_{\rm NL}^{(A)}\}$, $A=\rho,\, \mathcal{H},\, \mathcal{K}$ do not provide a complete scalar representation of the dynamics of LTB models, which is not surprising because the $\langle A\rangle_q[r_b]$ depend only on t and constitute boundary conditions for the A_q at $r=r_b$. From (39) and (61), local and non–local perturbations are related by

$$1+\delta_{\rm NL}^{(A)} = \frac{A_q(r)}{\langle A\rangle_q[r_b]}(1+\delta^{(A)}), \tag{63}$$

which shows that they only coincide at the boundary of each $\mathcal{D}[r_b]$ where $A_q(r_b) = \langle A\rangle_q[r_b]$. Since their time derivatives are evidently different, it is reasonable to expect that the evolution equations of the $\delta^{(A)}$ and $\delta_{\rm NL}^{(A)}$ will be different. Inserting (63) for $A=\rho,\, \mathcal{H}$ into (58c)–(58d) yields these equations:

$$\dot\delta_{\rm NL}^{(\rho)} = -\,3(1+\delta_{\rm NL}^{(\rho)})\,\langle\mathcal{H}\rangle_q[r_b]\,\delta_{\rm NL}^{(\mathcal{H})}, \tag{64a}$$

$$\begin{aligned}\dot\delta_{\rm NL}^{(\mathcal{H})} = &-(1+3\delta_{\rm NL}^{(\mathcal{H})})\,\langle\mathcal{H}\rangle_q[r_b]\,\delta_{\rm NL}^{(\mathcal{H})} + \frac{4\pi\,\langle\rho\rangle_q[r_b]}{3\langle\mathcal{H}\rangle_q^2[r_b]}(\delta_{\rm NL}^{(\mathcal{H})} - \delta_{\rm NL}^{(\rho)})\\ &-2\langle\mathcal{H}\rangle_q[r_b]\left(1-\frac{\mathcal{H}_q(r)}{\langle\mathcal{H}\rangle_q[r_b]}\right)^2 + 4(\mathcal{H}_q(r) - \langle\mathcal{H}\rangle_q[r_b])\delta_{\rm NL}^{(\mathcal{H})},\end{aligned} \tag{64b}$$

which, in order to render a fully complete system to describe the dynamics of the models in an arbitrary fixed domain $\mathcal{D}[r_b]$, needs to be supplemented by the evolution equations for $\langle\rho\rangle_q[r_b]$, $\langle\mathcal{H}\rangle_q[r_b]$ and ρ_q, $\mathcal{H}_q$:

$$\langle\mathcal{H}\rangle\dot{}_q[r_b] = -\langle\mathcal{H}\rangle_q^2[r_b] - \frac{4\pi}{3}\langle\rho\rangle_q[r_b], \qquad \dot{\mathcal{H}}_q(r) = -\mathcal{H}_q^2(r) - \frac{4\pi}{3}\rho_q(r), \tag{65a}$$

$$\langle\rho\rangle\dot{}_q[r_b] = -3\langle\mathcal{H}\rangle_q[r_b]\langle\rho\rangle_q[r_0], \qquad \dot{\rho}_q(r) = -3\mathcal{H}_q(r)\rho_q(r), \tag{65b}$$

while the constraints take the form (59) and with the $\delta^{(A)}$ expressed in terms of the $\delta^{(A)}_{\rm NL}$ by (63) (there is an extra constraint given by (17)). It is interesting to remark that (64*a*) is identical to (58*c*), but (64*b*) differs from (58*c*) by the terms with $\mathcal{H}_q$, which explains the need to add the evolution equations for ρ_q and $\mathcal{H}_q$. Notice that the non–local evolution equations become identical to the local ones (58*a*)–(58*d*) at the domain boundary in the limit $r \to r_b$ where $\mathcal{H}_q(r_b) = \langle\mathcal{H}\rangle_q[r_b]$. Also the non–local evolution equations depend on averages $\rho_q(r) = \langle\rho\rangle_q[r]$, $\mathcal{H}_q(r) = \langle\mathcal{H}\rangle_q[r]$ for inner points of $\mathcal{D}[r_b]$ and are considerably more complicated than (58*a*)–(58*d*). The self–consistency of the system (64*a*)–(65*b*) can be proved easily by comparing the mixed derivatives $[\dot{\delta}^{(A)}_{\rm NL}]'$ obtained from $[\delta^{(A)}_{\rm NL}]' = A'_q(r)/\langle A\rangle_q[r_b]$ with those that follow from the radial derivative of the right hand sides of (64*a*)–(64*b*).

8. Comparison with Buchert's averaging.

The most important difference between the q–average (13) and the standard proper volume average of Buchert's formalism (the average (13) with $\mathcal{F} = 1$) is the fact that the evolution equations that follow from (13) and its fluctuations lack the "back–reaction" correlation terms that appear in Buchert's evolution equations [41, 42, 43]. This fact follows readily by comparing (65*a*)–(65*b*) (or (58*a*)–((58*b*))) with the equivalent equations in Buchert's formalism.

Using (44) to express Σ in terms of $\mathcal{H} - \mathcal{H}_q = \mathbf{D}(\mathcal{H})$ and the commutation rule $\langle\mathcal{H}\rangle\dot{}_q - \langle\dot{\mathcal{H}}\rangle_q$ taken from (22), the weighed average (13) applied to the Raychaudhuri equation (8*a*) yields:

$$\langle\mathcal{H}\rangle\dot{}_q[r_b] = -\langle\mathcal{H}\rangle_q^2[r_b] - \frac{4\pi}{3}\langle\rho\rangle_q[r_b] + 2\mathcal{Q}_q[r_b], \tag{66}$$

where $\mathcal{Q}_q[r_b]$ is the back–reaction term:

$$\mathcal{Q}_q[r_b] \equiv \langle(\mathcal{H} - \langle\mathcal{H}\rangle_q)^2\rangle_q - \langle(\mathcal{H} - \mathcal{H}_q)^2\rangle_q = \langle[\mathbf{D}_{\rm NL}(\mathcal{H})\,]^2\rangle_q[r_b] - \langle[\mathbf{D}(\mathcal{H})]^2\rangle_q[r_b] = 0, \tag{67}$$

which vanishes identically for every domain as a consequence of (41) applied to $A = \mathcal{H}$ (see (C.1)–(C.3) for the proof of this result, see also [56]). Hence, (66) reduces exactly to the FLRW Raychaudhuri equation (65*a*) with zero back–reaction.

It is straightforward to prove that expressing the 1+3 local scalars (7) as (61) and (62) and applying the time derivative (22) to the Raychaudhury equation (8*a*) and to the remaining 1+3 equations (8*b*)–(8*d*), yields, for any domain $\mathcal{D}[r_b]$, the full set of evolution equations (64*a*)–(65*b*) without any back–reaction term (just as the substitution of (57*a*) and (57*b*) into (8*b*)–(8*d*) yields the system (58*a*)–((58*d*) without back–reaction terms). Hence, the weighed q–average (13), either with local or non–local fluctuations, leads to self–consistent and complete sets of evolution equations without back–reaction. This is a completely different outcome in comparison with the average from Buchert's formalism $\langle A\rangle_p[r_b]$ (notice the subindex $_p$). Applying Buchert's average to the Raychaudhuri equation (8a) yields a similar equation as (66), but with a non–vanishing back–reaction term [41, 42, 43]

$$\langle\mathcal{H}\rangle\dot{}_p = -\langle\mathcal{H}\rangle_p^2 - \frac{4\pi}{3}\langle\rho\rangle_p + 2\mathcal{Q}_p[r_b],$$
$$\text{with}: \qquad \mathcal{Q}_p[r_b] \equiv \langle(\mathcal{H}-\langle\mathcal{H}\rangle_p)^2\rangle_p - \langle(\mathcal{H}-\mathcal{H}_q)^2\rangle_p \neq 0, \qquad (68)$$

while the application of Buchert's average to the fluid flow equations (8c)–(8d) yields similar back–reaction terms (the energy balance equation (8b) remains with the same form). In particular, the Hamiltonian constraint (17) takes the form

$$\langle\mathcal{H}\rangle_p^2[r_b] = \frac{8\pi}{3}\langle\rho\rangle_p[r_b] - \langle\mathcal{K}\rangle_p[r_b] - \mathcal{Q}_p[r_b]. \qquad (69)$$

However, if we consider averaged scalars $\langle A\rangle_p$ and fluctuations constructed with this average in the framework of Buchert's formalism, we do not obtain a complete self–consistent set of evolution equations unless we make extra assumptions on the back–reaction terms [42, 43, 50].

Evidently, since invariant scalars like $\sigma_{ab}\sigma^{ab}$, $E_{ab}E^{ab}$ and $\sigma_{ab}E^{ab}$ are quadratic fluctuations with respect to the q–functions ρ_q, $\mathcal{H}_q$ (which relate to the q–averages $\langle\rho\rangle_q$ and $\langle\mathcal{H}\rangle_q$), and not with respect to functions associated with $\langle\rho\rangle_p$ and $\langle\mathcal{H}\rangle_p$, quadratic and higher order fluctuations in Buchert's formalism are not equal to these invariants, and thus we cannot obtain in this formalism the simple straightforward relations between averages of these invariants and statistical moments of ρ and $\mathcal{H}$ of section 6. However, we can still obtain analytic expressions for computing Buchert's average for these invariants. Considering that for any domain the q–average and Buchert's average of any scalar are related by

$$\langle A\rangle_q[r_b] = \frac{\langle A\,\mathcal{F}\rangle_p[r_b]}{\langle\mathcal{F}\rangle_p[r_b]}, \qquad (70)$$

together with the identities (52a)–(52c) and (53a)–(53c), we can obtain the following exact expressions for Buchert's average of quadratic scalar contractions:

$$\langle\sigma_{ab}\sigma^{ab}\rangle_p = \frac{\langle\mathcal{H}^2\,\mathcal{F}\rangle_p}{\langle\mathcal{F}\rangle_p} - \frac{\langle\mathcal{H}\,\mathcal{F}\rangle_p^2}{\langle\mathcal{F}\rangle_p^2} - \frac{\mathbf{Cov}_p(\sigma_{ab}\sigma^{ab},\,\mathcal{F})}{\langle\mathcal{F}\rangle_p}, \qquad (71a)$$

$$\langle E_{ab}E^{ab}\rangle_p = \frac{32\pi^2}{3}\left[\frac{\langle\rho^2\,\mathcal{F}\rangle_p}{\langle\mathcal{F}\rangle_p} - \frac{\langle\rho\,\mathcal{F}\rangle_p^2}{\langle\mathcal{F}\rangle_p^2}\right] - \frac{\mathbf{Cov}_p(E_{ab}E^{ab},\,\mathcal{F})}{\langle\mathcal{F}\rangle_p}, \qquad (71b)$$

$$\langle\sigma_{ab}E^{ab}\rangle_p = 8\pi\left[\frac{\langle\rho\,\mathcal{H}\,\mathcal{F}\rangle_p}{\langle\mathcal{F}\rangle_p} - \frac{\langle\rho\,\mathcal{F}\rangle_p\langle\mathcal{H}\,\mathcal{F}\rangle_p}{\langle\mathcal{F}\rangle_p^2}\right] - \frac{\mathbf{Cov}_p(\sigma_{ab}E^{ab},\,\mathcal{F})}{\langle\mathcal{F}\rangle_p}, \qquad (71c)$$

which lead to the following appealing identities which relate the two averages of these scalars and their correlation with the weight factor $\mathcal{F}$:

$$\langle\sigma_{ab}\sigma^{ab}\rangle_q - \langle\sigma_{ab}\sigma^{ab}\rangle_p = \frac{\mathbf{Cov}_p(\sigma_{ab}\sigma^{ab},\,\mathcal{F})}{\langle\mathcal{F}\rangle_p}, \qquad (72a)$$

$$\langle E_{ab}E^{ab}\rangle_q - \langle E_{ab}E^{ab}\rangle_p = \frac{\mathbf{Cov}_p(E_{ab}E^{ab},\,\mathcal{F})}{\langle\mathcal{F}\rangle_p}, \qquad (72b)$$

$$\langle\sigma_{ab}E^{ab}\rangle_q - \langle\sigma_{ab}E^{ab}\rangle_p = \frac{\mathbf{Cov}_p(\sigma_{ab}E^{ab},\,\mathcal{F})}{\langle\mathcal{F}\rangle_p}, \qquad (72c)$$

where we have omitted the $[r_b]$ symbol to simplify notation and $\mathbf{Cov}_p$ is the covariance correlation moment (42) with respect to Buchert's average.

9. A formalism of exact perturbations on a FLRW background.

It is evident that (as pointed out in previous work [35, 36]) the system (58a)–((58d) has the structure of evolution equations of spherical dust perturbations (the $\delta^{(A)}$) on a FLRW background (defined by the A_q). While the non–local perturbations were not considered in these references, the same resemblance to dust perturbation holds for systems like (64a)–(64b) and (65a)–(65b) involving $\langle A\rangle_q[r_b]$, $A_q(r)$ and $\delta^{(A)}_{\rm NL}(r, r_b)$.

Following the standard methodology [27, 28, 29, 63, 64], a perturbation formalism linking a "lumpy" spacetime (LTB model) and a homogeneous "background spacetime" (a FLRW dust model) in a domain $\mathcal{D}[r]$ can be defined by comparing local LTB variables with other LTB objects that define an FLRW "background state" of "zero order" variables by means of suitable maps. However, such maps must also deal with gauge issues involving a specific time slicing and coordinates. Since the boundary of every spherical comoving FLRW region can be mapped into the boundary of a domain $\mathcal{D}[r]$ of an LTB model by the Darmois matching conditions (31a)–(31b), and bearing in mind that LTB models and dust FLRW universes are both (i) spherically symmetric, (ii) have a geodesic 4–velocity and (iii) their full dynamics reduces to scalar modes [30, 32], a perturbation formalism associated with the q–scalars in domains $\mathcal{D}[r]$ can be defined rigorously by means of maps between covariant scalars with all gauge issues resolved.

9.1. The perturbation maps.

Let $\mathcal{M}$ be an LTB model and $X(\mathcal{D}[r])$ the set of all covariant scalars in an arbitrary domain $\mathcal{D}[r]$ of $\mathcal{M}$. Let $\tilde{\mathcal{M}}$ be a dust FLRW model and $\tilde X(\tilde{\mathcal{M}})$ the set of covariant scalars of $\tilde{\mathcal{M}}$:

> **The local perturbation map.** For every $\mathcal{D}[r]$ in $\mathcal{M}$ there exists a model $\tilde{\mathcal{M}}$ such that $A_q(t, r) = \tilde A(t)$ holds for every $\tilde A \in \tilde X(\tilde{\mathcal{M}})$ and $A \in X(\mathcal{M})$ and for all t. The following maps
>
> $$\Phi : \tilde X(\tilde{\mathcal{M}}) \to X(\mathcal{D}[r]), \qquad \tilde A \mapsto \Phi(\tilde A) = A_q(r) \in X(\mathcal{D}[r]), \tag{73}$$
>
> $$\delta^{(\ )} : X(\mathcal{D}[r]) \to X(\mathcal{D}[r]), \quad A \mapsto \delta^{(A)} = \frac{A - \Phi(\tilde A)}{\Phi(\tilde A)} = \frac{A - A_q}{A_q} = \frac{A - \langle A\rangle_q}{\langle A\rangle_q}, \tag{74}$$
>
> define for every $\mathcal{D}[r]$ a "background state" associated with an FLRW cosmology $\tilde{\mathcal{M}}$ and local exact perturbations of scalars A obtained by comparing them with the LTB scalars produced by the map Φ.

An analogous scalar perturbation formalism for the non–local fluctuations can be defined along the lines of (73) and (74):

> **The non–local perturbation map.** Let $Y(\mathcal{D}[r_b])$ be the set of all linear functionals in an arbitrary fixed comoving domain $\mathcal{D}[r_b] \subset {}^3\mathcal{T}[t]$. For every FLRW covariant scalar $\tilde A \in \tilde X(\tilde{\mathcal{M}})$ and every $\mathcal{D}[r_b]$ the following map
>
> $$\Phi_{\rm NL} : \tilde X(\tilde{\mathcal{M}}) \to Y(\mathcal{D}[r_b]), \qquad \tilde A \mapsto \langle A\rangle_q[r_b], \tag{75}$$
>
> defines a "background state" associated with a FLRW cosmology $\tilde{\mathcal{M}}$ but consisting of the functionals (13). The definition of the non–local

perturbation of the scalar A is then the map $\delta_{\rm NL}^{(\,)} : \tilde X(\tilde{\mathcal{M}}) \times X(\mathcal{D}[r_b]) \to X(\mathcal{D}[r_0]) \times Y(\mathcal{D}[r_0])$ such that

$$(\tilde A, A) \mapsto \delta_{\rm NL}^{(A)} = \frac{A - \Phi_{\rm NL}(\tilde A)}{\Phi_{\rm NL}(\tilde A)} = \frac{A(r) - \langle A\rangle_q[r_b]}{\langle A\rangle_q[r_b]}, \qquad (76)$$

which provides a pointwise comparison between the real valued functions A and their associated functionals $\langle A\rangle_q[r_b]$ assigned to the whole domain $\mathcal{D}[r_b]$.

It is important to emphasize that in the definitions above we distinguish between the FLRW "background state" and the FLRW "background spacetime" $\tilde{\mathcal{M}}$. As commented in section 4.4, the former is defined by the q–scalars: $\{\rho_q,\ \mathcal{H}_q\,\mathcal{K}_q,\ \Omega_q\}$ or $\{\langle\rho\rangle_q[r_b],\ \langle\mathcal{H}\rangle_q[r_b],\ \langle\mathcal{K}\rangle_q[r_b],\ \langle\Omega\rangle_q[r_b]\}$, which satisfy FLRW dynamics and relate to the FLRW scalars $\{\tilde\rho,\ \tilde{\mathcal{H}},\ \tilde{\mathcal{K}},\ \tilde\Omega\}$ of the background spacetime $\tilde{\mathcal{M}}$ by continuity under the Darmois matching conditions at an arbitrary constant r. These q–scalars are (in the perturbation maps) the "zero order" variables, with the "first order variables" being the local fluid flow scalars A related to the former by the perturbations $\delta^{(A)}$ or $\delta_{\rm NL}^{(A)}$.

Notice that the perturbation formalism defined by (73) and (74) is covariant because the background variables and the perturbations (zero and first order variables) are coordinate independent LTB objects (see sections 5 and 6). In fact, following the Stewart–Walker lemma [27, 28, 29, 65], the perturbations $\delta^{(A)}$ and $\delta_{\rm NL}^{(A)}$ are also gauge invariant (even in the usual sense of [66]) as they vanish in the FLRW spacetime $\tilde{\mathcal{M}}$ associated with the background state through (73) and (74).

9.2. LTB models as exact perturbations.

As opposed to the conventional approach to perturbations, either the traditional one with gauge invariant variables [63, 64, 66] or the covariant formalism of Ellis *et al* [27, 28, 29]), the perturbations that emerge from (73)–(74) and (75)–(76) are exact (not approximate) quantities, and thus do not lead to some unknown "near FLRW" space-time on the basis of a linearization process (though a rigorous linear limit can be defined, see section 12). Instead, the $\{A_q,\ \delta^{(A)}\}$ or the $\{\langle A\rangle_q[r_0],\ \delta_{\rm NL}^{(A)}\}$ with $A = \rho,\ \mathcal{H},\ \Omega,\ \mathcal{K}$ express a known class of spacetimes (generic LTB models) as exact spherical perturbations on an abstract FLRW background state defined by LTB objects: the A_q along all domains $\mathcal{D}[r]$ with varying boundary or the $\langle A\rangle_q[r_b]$ at a fixed domain. Evidently, these perturbation formalisms are special in the sense that they preserve the spherical symmetry and the dust source of the FLRW background. However, these formalisms are applicable to any space-time compatible with an LTB metric in the comoving frame and having an anisotropic fluid source [36], and can be readily generalized for the non–spherical Szekeres dust models [62].

10. Confined perturbations, Swiss cheese models and asymptotic perturbations.

In the definitions (73)–(74) and (75)–(76) we assumed domains $\mathcal{D}[r]$ or $\mathcal{D}[r_b]$ with r and r_b finite. As we show below, the resulting perturbations can also be considered in the asymptotic case when $r_b \to \infty$.



Figure 1. Local confined fluctuations. The fluctuations $\mathbf{D}(A)$ (thick vertical lines) follow by pointwise comparison of the scalars A and A_q along r in an arbitrary time slice ${}^3\mathcal{T}[t]$. Clump and void profiles are displayed in panels (a)–(b) and (c)–(d), the case of a Swiss cheese model matched to a FLRW region at $r = r_b$ is displayed in panels (b) and (d). Notice how in these latter cases $\mathbf{D}(A) = 0$ at $r = r_b$, thus forcing $A_q(r_b) = A(r_b)$, which explains the "humps" and "bags" in the radial profile of A.

10.1. Perturbations confined in comoving domains.

As long as r and r_b are finite, we are considering perturbations that are confined in arbitrary bounded comoving domains of an LTB model. Depending on whether we perform a smooth match with a FLRW spacetime or not we have the following possibilities:

- **There is no matching with FLRW** (see panels (a) and (c) of figures 1, 2 and 3). The perturbation describes the dynamics of the domain $\mathcal{D}[r]$ or $\mathcal{D}[r_b]$ with respect to a FLRW background state defined by (73) or (75), which is a fictitious reference FLRW dust model $\tilde{\mathcal{M}}$ (the background spacetime) whose scalars $\tilde{A}$ match (via Darmois matching conditions) the zero order variables $A_q(r)$ or $\langle A\rangle_q[r_b]$. Notice that:
  - The fictitious reference FLRW dust model $\tilde{\mathcal{M}}$ is necessarily different for different domains.
  - First order quantities like A and the gradients A' and A'_q ($A = \rho$, $\mathcal{H}$, $\mathcal{K}$, Ω) are not continuous at the domain boundaries r or r_b. See section 4.4 and



(a) (b) (c) (d)

Figure 2. Non–local confined fluctuations. The panels display (as in figure 1) clump and void profiles with and without a Swiss cheese matching to a FLRW region at $r = r_b$ in an arbitrary time slice ${}^3\mathcal{T}[t]$. The non–local fluctuations $\mathbf{D}_{\mathrm{NL}}(A)$ (the arrows) compare local values of $A(r)$ with the average $\langle A\rangle_q[r_b]$ (thick line) assigned to the whole domain $\mathcal{D}[r_b]$ (shaded region). Notice (as in figure 1) that $\mathbf{D}_{\mathrm{NL}}(A)$ only vanishes at $r = r_b$ if there is a Swiss cheese matching (panels (b) and (d)).

panels (a) and (c) of figures 1, 2 and 3.

- **Swiss cheese holes smoothly matched to a FLRW region** (see panels (b) and (d) of figures 1, 2 and 3). The perturbation describes the dynamics of a comoving domain $\mathcal{D}[r_b]$ with respect to a background state that in this case corresponds to the actual (non–fictitious) FLRW background spacetime matched at $r = r_b$ (under conditions (31a)–(31c)). Notice that:
  - The resulting spacetime is a compound Swiss cheese configuration consisting of an LTB section (confined in $\mathcal{D}[r_b]$) described by zero order variables A_q and their perturbations $\delta^{(A)}$ or $\delta^{(A)}_{\mathrm{NL}}$, and the FLRW background spacetime that extends for $r > r_b$. The FLRW dust model $\tilde{\mathcal{M}}$ is necessarily different for different domains $\mathcal{D}[r_b]$.
  - Darmois matching conditions only require the zero order quantities A_q and $\langle A\rangle_q[r_b]$ to be continuous at r_b, though it is always possible to demand (as extra conditions) that the local scalars A and the gradients A' and A'_q (which are first order quantities related to $\delta^{(A)}$ via (35)) be also continuous at

r_b. These extra conditions imply that the fluctuations and perturbations themselves vanish at r_b (see section 4.4 and panels b and d of figures 1, 2 and 3).

10.2. Asymptotic perturbations.

The application of the perturbation maps (73)–(74) and (75)–(76) to the asymptotic limit $r \to \infty$ or $r_b \to \infty$ depends on the convergence of LTB scalars and their associated q–scalars along radial rays (*i.e.* curves $x^a = \ell\delta^a_r$, where $\ell = \int g_{rr}\mathrm{d}r$ is the proper radial length), which are spacelike geodesics of the metric (1) and of the slices ${}^3\mathcal{T}[t]$ with metric h_{ab}. As shown in [39], $\ell(r)$ is a positive monotonic function in regular LTB models, and thus the asymptotic limit $r \to \infty$ corresponds to $\ell \to \infty$.

The radial asymptotic behavior of LTB models is determined by the following asymptotic limits for the scalars $A = \rho$, $\mathcal{H}$, $\mathcal{K}$, Ω: [+]

$$\lim_{r\to\infty} A(r) = \lim_{r\to\infty} A_q(r) = \lim_{r_b\to\infty} \langle A\rangle_q[r_b] = \begin{cases} A_\infty(t) \neq 0, & \text{Asymptotically FLRW} \\ \\ 0, & \text{Asymptotically Minkowski.} \end{cases} \tag{77}$$

These limits correspond to q–averages for increasingly large domains up to the situation in which $\mathcal{D}[r]$ or $\mathcal{D}[r_b]$ become the whole slice ${}^3\mathcal{T}[t]$, and thus, we can speak of the q–average of a whole LTB model (instead of the q–average of confined domains of an LTB model). We look at the perturbations for asymptotically FLRW and Minkowski models separately below.

- **Asymptotically FLRW models** (see figure 4). As a consequence of (74) and (77) local perturbations vanish asymptotically in these models:

$$\lim_{r\to\infty} \delta^{(A)}(r) = 0. \tag{78}$$

  However, (76) and (77) lead for non–local perturbations to the following non–trivial nonzero asymptotic limit:

$$\lim_{r_b\to\infty} \delta^{(A)}_{\text{NL}}(r, r_b) = \frac{A(r) - A_\infty(t)}{A_\infty(t)} = \frac{A(r) - \tilde{A}(t)}{\tilde{A}(t)}, \qquad r \text{ finite}, \tag{79}$$

  which keeps their interpretation of a "contrast" between local values $A(r)$ and the domain average, with the difference that the average now corresponds to a domain that encompasses the whole slice ${}^3\mathcal{T}[t]$ and coincides with the corresponding scalars $A_\infty(t) = \tilde{A}(t)$ of an asymptotic FLRW background state defined by (75). As a consequence, we can rigorously state that the q–average of every LTB model of this class is exactly the FLRW background spacetime $\tilde{\mathcal{M}}$. Asymptotic fluctuations and perturbations for a model converging to FLRW are illustrated by figure 4.

- **Asymptotically Minkowski models**. By looking at (76) it is evident that non–local perturbations $\delta^{(A)}_{\text{NL}}$ cannot be defined in the asymptotic limit for these LTB models, since in general $A(r) \neq 0$ for finite r and thus the ratio $A/\langle A\rangle_q[r_b]$ diverges as $\langle A\rangle_q[r_b] \to 0$ in the limit $r_b \to \infty$. However, local perturbations $\delta^{(A)}$ defined by (74) are not affected by the radial convergence to a Minkowski vacuum

---

[+] LTB models can only converge in the asymptotic radial regime to FLRW dust models with zero or negative spatial curvature, not positive. Models convergent to Minkowski can converge to a section of Minkowski parametrized by Milne coordinates or by non–standard curvilinear coordinates. In the Milne case ρ_q, $\Omega_q \to 0$ but $\mathcal{H}_q$, $\mathcal{K}_q$ tend to nonzero values (see [39]).

because, as shown in [39], both $A \to 0$ and $A_q \to 0$ hold as $r \to \infty$ but their ratio A/A_q is finite in this limit, and thus the $\delta^{(A)}$ in (74) are well defined. The only difference with asymptotically FLRW models is that the $\delta^{(A)}$ tend (in general) to nonzero constant values as $r \to \infty$ (see [39]).

11. Comparison between local and non–local perturbations.

Since the perturbations $\delta^{(A)}$ and the $\delta^{(A)}_{\mathrm{NL}}$ are different objects, they provide a different measure of the inhomogeneity or "deviation" of an LTB model from a FLRW Universe:

- The $\delta^{(A)}$ measure this deviation through local comparison of $A(r)$ with the characteristic FLRW scalar $A_q(r) = \langle A\rangle_q[r]$ at every r, and is related (via (35)) to radial gradients of A_q and A and (via (49a)–(49b)) to the ratio of Weyl to Ricci curvature and anisotropic to isotropic expansion. See figures 1 and 3.
- The $\delta^{(A)}_{\mathrm{NL}}$ measure the deviation from FLRW though the notion of the "contrast" between local values of $A(r)$ inside an arbitrary but fixed comoving domain ($0 \le r < r_b$) and a value $\langle A\rangle_q[r_b]$ that can be associated to the equivalent FLRW scalar $\tilde{A}(t) = \langle A\rangle_q[r_b]$ that characterizes the whole domain. In the asymptotic limit $r_b \to \infty$ so that $\tilde{A}(t) = A_\infty(t)$, these perturbations provide the contrast with respect to the FLRW spacetime to which the LTB model converges asymptotically. See figures 2 and 3.

We examine below how these differences relate to the shape of the radial profile of a scalar at an arbitrary ${}^3\mathcal{T}[t]$ and the signs and magnitudes ("amplitude") of the perturbations.

11.1. Signs of the perturbations vs type of radial profiles.

Consider a scalar A whose radial profile is monotonic throughout a given domain *. Then, following (23), the clump/void profiles are defined by

$$\text{Clump profile} \qquad A' \le 0 \quad \Rightarrow \quad A'_q \le 0 \quad \Rightarrow \quad A \le A_q, \tag{80a}$$

$$\text{Void profile} \qquad A' \ge 0 \quad \Rightarrow \quad A'_q \ge 0 \quad \Rightarrow \quad A \ge A_q, \tag{80b}$$

where $A(0) = A_q(0) = A_c$ and $A'(0) = A'_q(0) = 0$ and we have assumed absence of shell crossings and singular layers (hence $R' > 0$ holds everywhere or it has a common same order zero with A' and A'_q [39, 40]). The relation between the signs of the perturbations and the profile type follows from (35) (it can also be seen in figures 1, 2 and 3):

- **Local perturbations**:

$$\mathrm{sign}(\delta^{(A)}) = \mathrm{sign}(A'_q) \quad \Rightarrow \quad \left\{ \begin{array}{ll} \text{clump profile}: & \delta^{(A)} \le 0, \\ & \\ \text{void profile}: & \delta^{(A)} \ge 0. \end{array} \right. \tag{81}$$

---

* Such domains always exist for r_b finite, even if A is not monotonic for values $r > r_b$. The nature and evolution of radial profiles of LTB scalars were examined extensively in reference [40]. In general, density void profiles are fully compatible with regularity conditions (absence of shell crossings) for hyperbolic models, but not for elliptic or parabolic ones. Also, for models with a non–simultaneous big bang ($t'_{\mathrm{bb}} \ne 0$, nonzero decaying modes) density void profiles always emerge from initial clump profiles after a transition ("profile inversion"), whereas a density void profile exists for all the time evolution only in models with a simultaneous bang (zero decaying modes). A profile inversion of the expansion scalar $\mathcal{H}$ necessarily occurs in elliptic models but not in hyperbolic ones (see [40] for further detail).

- **Non–local perturbations**:
  Finite domains $\mathcal{D}[r_b]$ and Swiss cheese holes. The sign relations in (80a) and (80b) imply that there always exists a value $\bar r = y < r_b$ such that $A(y) = A_q(r_b) = \langle A\rangle_q[r_b]$, and thus we have from (39): $\delta^{(A)}_{\rm NL}(y, r_b) = 0$. Since $A_c > \langle A\rangle_q[r_b]$ holds in clump profiles and $A_c < \langle A\rangle_q[r_b]$ in void profiles, then $\delta^{(A)}_{\rm NL}$ changes sign as follows (see panels (b) and (d) of figures 2 and 3):

  $$\text{Clump:}\qquad \delta^{(A)}_{\rm NL} > 0 \;\text{ for }\; 0 \le \bar r < y, \qquad \delta^{(A)}_{\rm NL} < 0 \;\text{ for }\; y < \bar r < r_b, \tag{82a}$$

  $$\text{Void:}\qquad \delta^{(A)}_{\rm NL} < 0 \;\text{ for }\; 0 \le \bar r < y, \qquad \delta^{(A)}_{\rm NL} > 0 \;\text{ for }\; y < \bar r < r_b, \tag{82b}$$

  Asymptotic background (see figure 4). In the limit $r_b \to \infty$ we have $A \to A_{\infty}$, $\langle A\rangle_q[r_b] \to A_{\infty}$ and $\mathcal{D}[r_b] \to {}^3\mathcal{T}[t]$, hence for all r:

  $$\text{Clump:}\qquad A > A_{\infty} \qquad \delta^{(A)}_{\rm NL} > 0, \tag{83a}$$

  $$\text{Void:}\qquad A < A_{\infty} \qquad \delta^{(A)}_{\rm NL} < 0, \tag{83b}$$

Comparing the sign relations in (81) with those in (82a)–(82b) and (83a)–(83b) clearly indicates that non–local perturbations $\delta^{(A)}_{\rm NL}$ have the expected intuitive signs that we tend to infer for perturbations through the familiar notion of a "density contrast" with respect to a fiducial (or background) FLRW density value $\tilde\rho$: an over–density (clump) is a positive perturbation because local density is larger than $\tilde\rho$, while an under–density (void) is a negative perturbation because $\tilde\rho$ is smaller than local density, which we can now identify with $\langle\rho\rangle_q[r_b]$ or its asymptotic limit ρ_{∞}. In fact, as shown in figures 2, 3 and 4, this intuitive notion of a "contrast" can be applied to all covariant scalars, not just the density. On the other hand, the "gradient" based local perturbations $\delta^{(A)}$ have the opposite sign to the intuitively expected one and thus are more abstract and counter–intuitive (see figures 3 and 4).

11.2. Amplitudes.

The difference between the perturbations $\delta^{(A)}$ and $\delta^{(A)}_{\rm NL}$ can also be understood in terms of the notion of "amplitude" often introduced in the context of simple intuitive Newtonian [67] and relativistic [3, 4, 10] perturbations. In order to illustrate this point we consider an asymptotic perturbation in an LTB model converging radially to FLRW (as in figure 4), so that for all A we have along the slices ${}^3\mathcal{T}[t]$ the asymptotically FLRW behavior of (77):

$$\lim_{r_b\to\infty} A(r_b) = \lim_{r_b\to\infty} \langle A\rangle_q[r_b] = A_{\infty}(t) = \tilde A(t) \neq 0, \tag{84}$$

where $\tilde A(t)$ is the A scalar for the asymptotic FLRW state (see figure 4). Assuming for simplicity a monotonic profile, then without loosing generality we can express any A in any complete slice ${}^3\mathcal{T}[t]$ (not intersecting a singularity) in terms of a "contrast amplitude" ϕ_A as

$$A = A_{\infty}\left[1 + \phi_{A}\, f(r)\right], \quad \text{with} \quad \phi_{A} \equiv \delta^{(A)}_{\rm NL}(0,\infty) = \lim_{r_b\to\infty}\left[\frac{A_c}{\langle A\rangle_q[r_b]} - 1\right] = \frac{A_c}{A_{\infty}} - 1, \tag{85}$$

where $A_c = A(t, 0) = A_q(t, 0)$ and f is a smooth non–negative function satisfying $f_c = f(0) = 1$ and $f'(0) = 0$, as well as $f(r)$, $f'(r) \to 0$ as $r \to \infty$. Since we can



(a) (b)

(c) (d)

Figure 3. Confined perturbations. The panels display the local and non–local perturbations ($\delta^{(A)}$ and $\delta^{(A)}_{\rm NL}$) for the radial profiles displayed in figures 1 and 2. Notice that both types of perturbations vanish at $r = r_b$ when there is a Swiss cheese matching (panels (b) and (d)). Notice also that $\delta^{(A)}$ and $\delta^{(A)}_{\rm NL}$ have opposite signs around the center.

always choose at any ${}^3\mathcal{T}[t]$ the radial coordinate as $R = r$, so that $R^2R'\mathrm{d}r = r^2\mathrm{d}r$, the form (85) implies from (13) and (18)

$$A_q(r) = A_\infty\left[1 + \phi_A f_q(r)\right], \qquad f_q(r) = \langle f\rangle_q[r] = \frac{\int_0^r f\,\bar r^2\mathrm{d}\bar r}{\int_0^r \bar r^2\mathrm{d}\bar r}, \tag{86}$$

while (34), (39) and (63) yield

$$\delta^{(A)}_{\rm NL} = \phi_{_A}\, f \qquad \delta^{(A)} = \frac{(f - f_q)\phi_{_A}}{1 + \phi_A\, f_q}, \tag{87}$$

which illustrate the following basic features:

- The contrast amplitude $\phi_{_A}$ obeys a simple linear relation with the more intuitive non–local perturbations $\delta^{(A)}_{\rm NL}$, but its relation with local perturbations $\delta^{(A)}$ is non–linear. This is consistent with the fact that the latter perturbations are less intuitive than the former.
- Both perturbations vanish in the domain boundary $r \to \infty$ where the correspondence with the FLRW background occurs, but their behavior at the center is different: $\delta^{(A)}_{\rm NL}(0,\infty) = \phi_{_A} \neq 0$ (in general) while $\delta^{(A)}(0) = 0$ (see figure 4).



Figure 4. Asymptotic fluctuations and perturbations. Panels (a) and (b) correspond to a clump profile, while (c) and (d) depict a void profile. Since the domain encompasses a whole time slice ${}^3\mathcal{T}[t]$, non–local fluctuations compare local values of $A(r)$ with the global average $\langle A\rangle_q[r_b] \to A_\infty$ in the limit $r_b \to \infty$, which is equal to the corresponding scalar $\tilde{A}(t)$ of the FLRW asymptotic state. Non–local perturbations $\delta^{(A)}_{\rm NL}$ yield the "contrast" between local values of A and this average. The amplitude of the perturbation is ϕ_A, while ψ_A depicts the maximal value of the local perturbation $\delta^{(A)}$. Notice that $\delta^{(A)}_{\rm NL}$ is positive/negative for the clump/void, while $\delta^{(A)}$ has opposite signs. Also, both perturbations vanish as $r \to \infty$.

- The relation between the signs of the perturbations and the type of profile can also be given in terms of the amplitude:

$$\text{Clump profile:}\quad A_c > A_\infty,\quad \phi_A > 0,\qquad \delta^{(A)}_{\rm NL} > 0 \quad\text{but}\quad \delta^{(A)} < 0, \tag{88a}$$

$$\text{Void profile:}\quad A_c < A_\infty,\quad \phi_A < 0,\qquad \delta^{(A)}_{\rm NL} < 0 \quad\text{but}\quad \delta^{(A)} > 0. \tag{88b}$$

  where we used the fact that (by construction) $f' \leq 0$ so that the sign relations (80a)–(80b) imply $f - f_q \leq 0$ and the sign of $\delta^{(A)}$ is the opposite of the sign of ϕ_A.

Evidently, the convention whereby a clump or void (*i.e.* overdensity or underdensity when $A = \rho$) respectively correspond to positive or negative contrast follows from the sign of the amplitude contrast embodied in $\delta^{(A)}_{\rm NL}$, and is more intuitive and easier to follow than the complicated relation between radial gradients A'_q associated with $\delta^{(A)}$.

However, in spite of their differences in signs when referred to clump/void radial profiles, the deviation form homogeneity (FLRW conditions) can be traced effectively with both types of perturbations, $\delta^{(A)}_{\rm NL}$ and $\delta^{(A)}$, because there is a consistent



Figure 5. Magnitudes of local and non–local perturbations. The graph depicts the monotonic non–linear relation between the amplitude, ϕ_A, of non–local perturbations and the maximal value of the local perturbation, ψ_A (see figure 4), which shows that both perturbations provide a self–consistent qualitatively analogous description of the deviation from FLRW. Only for small amplitudes $|\phi_A| \ll 1$ the relation is linear.

monotonic relation between their magnitudes (their maximal/minimal value in a given domain) and the amplitude ϕ_A. For the non–local perturbations $\delta_{\rm NL}^{(A)}$ the amplitude $\phi_{_A}$ coincides with their extremal value, which occurs at the center $r = 0$ (for domains with monotonic f. For local perturbations the relation between ϕ_A and the extrema of $\delta^{(A)}$ (denoted by $\psi_{_A}$, see figure 4) is more complicated, as the latter may occur for different values of r for different slices ${}^3\mathcal{T}[t]$ and depends on the choice of f. Since a result for a general f is hard to find, we consider the special form $f = (1 + r^n)^{-1}$ where $n > 0$ (the results are qualitatively analogous for all other forms). The result, shown in figure 5, illustrates the non–linear monotonic relation between $\psi_{_A}$ and ϕ_A. Hence, both types of perturbations provide a consistent estimate of the deviation from homogeneity of the models in which larger values of $|\phi_A|$ correspond to larger magnitudes of $\delta^{(A)}$ and $\delta_{\rm NL}^{(A)}$.

12. Linear limit.

While the $\delta^{(A)}$ and the $\delta_{\rm NL}^{(A)}$ are exact quantities that need not be "small' and do not comply with linear evolution equations, we can expect (intuitively) that they should somehow reduce to linear dust perturbations (in the comoving gauge) when their magnitudes (*i.e.* amplitudes) are small. A more rigorous way to examine their connection to linear perturbations of dust sources follows by constructing second order equations for $\delta^{(\rho)}$ and $\delta_{\rm NL}^{(\rho)}$:

- **Local perturbations**. We differentiate both sides of (58*a*) and use the remaining equations (58*b*)–(58*d*) to eliminate all derivatives except $\ddot{\delta}^{(\rho)}$ and $\dot{\delta}^{(\rho)}$, leading to:

$$\ddot{\delta}^{(\rho)} - \frac{2\,[\dot{\delta}^{(\rho)}]^2}{1+\delta^{(\rho)}} + 2\mathcal{H}_q\,\dot{\delta}^{(\rho)} - 4\pi\rho_q\,\delta^{(\rho)}\left(1+\delta^{(\rho)}\right) = 0, \qquad (89)$$

- **Non–local perturbations**. We follow the same procedure as above: differentiate (64*a*) and use (64*b*)–(65*b*) to rewrite it in factors of $\dot{\delta}_{\rm NL}^{(A)}$ and $\delta_{\rm NL}^{(A)}$.

Considering asymptotic backgrounds ($r_b \to \infty$) yields:

$$\ddot{\delta}^{(\rho)}_{\text{NL}} - \frac{2\,[\dot{\delta}^{(\rho)}_{\text{NL}}]^2}{1+\delta^{(\rho)}_{\text{NL}}} + 4\,(6\mathcal{H}_\infty - 4\mathcal{H}_q)\,\dot{\delta}^{(\rho)}_{\text{NL}} - \left[4\pi\rho_\infty\,\delta^{(\rho)}_{\text{NL}} + 6\,(\mathcal{H}_q - \mathcal{H}_\infty)^2\right]\left(1+\delta^{(\rho)}_{\text{NL}}\right) = 0, \tag{90}$$

which holds for confined domains by replacing ρ_∞, $\mathcal{H}_\infty$ with $\langle\rho\rangle_q[r_b]$, $\langle\mathcal{H}\rangle_q[r_b]$.

We remark that both (89) and (90) are exact non–linear equations for $\delta^{(\rho)}$ or $\delta^{(\rho)}_{\text{NL}}$ (equations similar to (89) have been obtained in [68, 69] for Szekeres models). Also, notice that (89) and (90) coincide as $r \to \infty$ for asymptotic perturbations and as $r \to r_b$ for confined domains $\mathcal{D}[r_b]$.

In general there is no reason to assume that the amplitudes ϕ_A or their time derivatives $\dot{\phi}_A$ are small in a generic LTB model subjected to non–linear evolution equations, but if in a given range of t we have $|\phi_A| \ll 1$ then the relation between both $\delta^{(A)}$ and $\delta^{(A)}_{\text{NL}}$ with ϕ_A should become similar. This can be seen by expanding for $|\phi_A| \ll 1$ in the non–linear relation in (87) between the $\delta^{(A)}$ and the amplitude ϕ_A:

$$\delta^{(A)} \approx (f - f_q)\,\phi_A + O(\phi_A^2), \tag{91}$$

so that the $\delta^{(A)}$ become linear on ϕ_A (as the $\delta^{(A)}_{\text{NL}}$), and thus the magnitudes of $\delta^{(A)}$ and $\delta^{(A)}_{\text{NL}}$ are both proportional to ϕ_A because $0 < f \leq 1$ and the fluctuations $f - f_q$ are bounded. As a consequence, if $|\phi_A| \ll 1$ both perturbations are of the same order in $\phi_{_A}$, and thus both comply with $|\delta^{(A)}| \ll 1$ and $|\delta^{(A)}_{\text{NL}}| \ll 1$ and the time evolution of $\delta^{(A)}$ must be the same as that of $\delta^{(A)}_{\text{NL}}$ at leading order in $\phi_{_A}$. In particular, if we take $A = \rho, \mathcal{H}$, the linearity conditions $|\phi_{_\rho}| \ll 1$ and $|\phi_{_\mathcal{H}}| \ll 1$ imply that $|\delta^{(\rho)}| \ll 1$, $|\delta^{(\rho)}_{\text{NL}}| \ll 1$ and $|\delta^{(\mathcal{H})}| \ll 1$, $|\delta^{(\mathcal{H})}_{\text{NL}}| \ll 1$ hold, and thus $|\dot{\delta}^{(\rho)}| \ll 1$, $|\dot{\delta}^{(\rho)}_{\text{NL}}| \ll 1$ also holds (from (58*c*) and (64*a*)). Therefore, we also have $\rho_q \approx \rho_\infty \approx \rho$ and $\mathcal{H}_q \approx \mathcal{H}_\infty \approx \mathcal{H}$ (the same relations hold for finite domains by replacing ρ_∞, $\mathcal{H}_\infty$ with $\langle\rho\rangle_q[r_0]$, $\langle\mathcal{H}\rangle_q[r_0]$). Considering all these implications, if $|\phi_A| \ll 1$ the second order evolution equations (89) and (90) becomes at order $\phi_{_A}$

$$\ddot{\delta}^{(\rho)} + 2\mathcal{H}_\infty\,\dot{\delta}^{(\rho)} - 4\pi\rho_\infty = 0, \tag{92}$$

which is formally identical to the evolution equation for gauge invariant linear density perturbations of a dust source around a FLRW background characterized by ρ_∞, $\mathcal{H}_\infty$ (or $\langle\rho\rangle_q[r_b]$, $\langle\mathcal{H}\rangle_q[r_b]$ for bounded domains) in the comoving gauge [67, 70], which for dust is a synchronous gauge as well.

13. Gravitational entropy.

A gravitational entropy functional has been proposed by Buchert *et al.* [42, 57, 58, 59], as an application of the Kullback–Leibler functional of Information Theory to spacetimes with inhomogeneous dust sources. While Buchert and coworkers defined this functional with the standard proper volume average (*i.e.* (13) with $\mathcal{F} = 1$), as this is the scalar average of Buchert's formalism, we consider here the same functional with the weighed q–average (13) with $\mathcal{F} \neq 1$. Hence, our approach will yield different theoretical connections.

Following Buchert *et al*, we can define the Kullback–Leibler entropy functional in terms of the q–average over a domain $\mathcal{D}[r_0]$ as

$$S - S_{\text{eq}} = \gamma_0 \int_{\mathcal{D}[r_b]} \rho\,\ln\left[\frac{\rho}{\langle\rho\rangle_q[r_b]}\right]\mathcal{F}\mathrm{d}\mathcal{V}_p = \gamma_0\left\langle \rho\ln\frac{\rho}{\langle\rho\rangle_q[r_b]}\right\rangle_q [r_b]\,\mathcal{V}_q(r_b), \tag{93}$$

$$\text{with :}\qquad \mathcal{V}_q = \int_{\mathcal{D}[r_b]} \mathcal{F}\,\mathrm{d}\mathcal{V}_p = \frac{4\pi}{3}R^3(r_b), \tag{94}$$

where $S_{\text{eq}} = S_{\text{eq}}(r_b)$, and we have introduced the constant $\gamma_0 = k_{_B}/(mc^2)$, with $k_{_B}$ being the Boltzmann's and m a particle mass, so that $S - S_{\text{eq}}$ has units of entropy. If we define the entropy current associated with (93) as $S^a = \rho\,(S - S_{\text{eq}})\,u^a$, then by considering the rest mass density conservation law (8*b*) we obtain the following entropy balance law:

$$\nabla_a S^a \geq 0 \qquad\Rightarrow\qquad \dot{S} \geq 0, \tag{95}$$

which allows us to identify the inhomogeneity measured by $\rho(r) \neq \langle\rho\rangle_q[r_b]$ as the source of gravitational entropy production ($\dot{S} > 0$) in arbitrary domains $\mathcal{D}[r_b]$ of LTB models, with equilibrium states complying with $\dot{S} = 0$, $S = S_{\text{eq}}$ and corresponding to:

- FLRW models as the homogeneous subset (particular case) of LTB models for which $\rho(r) = \langle\rho\rangle_q[r_b]$ holds for all r, r_b, t.
- specific boundaries of generic LTB models, such as:
    - The center worldline (in models admitting a regular center) corresponding to the case $r_b = 0$, since $\rho(t,0) = \rho_q(t,0)$ holds for all t [39, 40, 56].
    - Asymptotic boundary $r \to \infty$ along radial rays of time slices ${}^3\mathcal{T}[t]$ of LTB models radially converging to a FLRW state (see section 10.2 and [39]), for which both ρ and ρ_q tend to the same value $\rho_{_\infty} = \tilde{\rho}(t)$ in this limit (see figure 4).
    - The boundary $r = r_b$ (for all t) of Swiss cheese holes in domains $\mathcal{D}[r_b]$ smoothly matched with a FLRW region when the extra conditions (32) are imposed, so that $\rho(r_b) = \rho_q(r_b)$ holds (see sections 4.4 and 10.1, and panels (b) and (d) of figures 1, 2 and 3). Evidently $\dot{S} = 0$ holds for all t in the FLRW region $r > r_b$.

We remark that in all the equilibrium states listed above we have $\delta^{(\rho)} = \delta^{(\rho)}_{\text{NL}} = 0$, and thus entropy production is closely related with nonzero density perturbations. For the remaining of this section we will assume domains such that $\dot{S} \neq 0$ holds.

In order to evaluate $\dot{S}$, we apply the time derivative commutation rule (22) to (93), leading after some algebraic manipulation to the same relation between $\dot{S}$ and the non–commutativity of the time derivative and the average found in [57, 58, 59]:

$$\frac{\dot{S}(r_b)}{\gamma_0\mathcal{V}_q(r_b)} = \langle\dot{\rho}\rangle_q[r_b] - \langle\rho\rangle\dot{}_q[r_b], \tag{96}$$

which, with the help of (22) and (42), can be related to the negative statistical correlation of fluctuations of ρ and $\mathcal{H}$, given by the covariance momentum with respect to the q–averages of these variables in a domain $\mathcal{D}[r_b]$:

$$\frac{\dot{S}}{\mathcal{V}_q} = -3\gamma_0\left[\,\langle\rho\mathcal{H}\rangle_q - \langle\rho\rangle_q\langle\mathcal{H}\rangle_q\,\right] = -3\gamma_0\mathbf{Cov}_q(\rho,\mathcal{H}) = -3\gamma_0\langle\mathbf{D}(\rho)\,\mathbf{D}(\mathcal{H})\rangle_q \geq 0, \tag{97}$$

so that:

$$\mathbf{Cov}_q(\rho,\mathcal{H}) = \langle\mathbf{D}(\mathcal{H})\mathbf{D}(\rho)\rangle_q[r_b] < 0 \;\Rightarrow\; \dot{S}(r_b) > 0, \tag{98}$$

where we have applied (42) to replace non–local fluctuations with local ones (see proof in Appendix C). Considering (5)–(6), (44), (46*c*) and (52*c*), condition (98) can be given in terms of the q–average of a scalar invariant by:

$$\langle\sigma_{ab}\,E^{ab}\rangle_q[r_b] < 0 \;\Rightarrow\; \dot{S}(r_b) > 0, \tag{99}$$

which is a very elegant way to connect (93) with an unequivocal and completely coordinate independent marker of inhomogeneity, as it contains contributions from density and velocity fluctuations. Notice that, as consequence of (97)–(99), the condition $\dot S = 0$ in equilibrium states implies $\delta^{(\rho)} = \delta^{(\rho)}_{\rm NL} = \delta^{(\mathcal{H})} = \delta^{(\mathcal{H})}_{\rm NL} = 0$, and thus perturbations of all q–scalars necessarily vanish in these states.

As pointed out by Buchert *et al* in [57, 58, 59] (who also obtained the relations (97) and (98) but not (99)), a negative correlation of the density and Hubble fluctuations seems to be consistent with the intuitive behavior of gravitational clustering processes in structure formation scenarios: over/under dense regions tend to contract/expand as density increases/decreases. However, this intuitive behavior occurs in the time evolution, not (necessarily) in domains along time slices where averages are computed. Therefore, as much as (98) is an intuitively appealing and elegant proposition (in view of (99)), it is not *a priori* evident that we will obtain the desired negative sign of $\mathbf{D}(\rho)\mathbf{D}(\mathcal{H})$ when actually computing the fluctuations. Since verifying (98) in arbitrary domains of generic inhomogeneous LTB models is a comprehensive task that is beyond the scope of this paper, and thus will be accomplished in a future separate article [71], we provide in this section a guideline on how this verification can be accomplished, as well as some preliminary results (listed in the following section).

Testing (98) involves computing radial integrals of quantities whose explicit radial dependence is not known (and thus must be done numerically). However, we can infer the fulfillment of (98) by means of the following sufficient (but not necessary) condition that can be evaluated at the boundary of every domain $r = r_0$:

$$\mathbf{D}(\mathcal{H})\mathbf{D}(\rho) = [\,\mathcal{H}(r_b) - \mathcal{H}_q(r_b)\,][\,\rho(r_b) - \rho_q(r_b)\,] < 0 \;\Rightarrow\; \dot S(r_b) > 0, \tag{100}$$

where we used (42) (see its proof in (C.6)–(C.7)) to be able to consider the fluctuations evaluated at the boundary $r = r_b$ of a fixed but arbitrary domain. Considering (23) and (24), the product of these fluctuations is

$$\mathbf{D}(\mathcal{H})\mathbf{D}(\rho) = \frac{\rho_q'(r_b)\,\mathcal{H}_q'(r_b)}{[\mathcal{V}_q'(r_b)/\mathcal{V}_q'(r_b)]^2} = \frac{1}{\mathcal{V}_q^2(r_b)}\int_0^{r_b}\rho'\,\mathcal{V}_q\,\mathrm{d}\bar r\int_0^{r_b}\mathcal{H}'\,\mathcal{V}_q\,\mathrm{d}\bar r, \tag{101}$$

and thus condition (100) can be given directly in terms of the correlation of the radial gradients of the density and the Hubble expansion scalar: ♯

$$\rho_q'(r_b)\,\mathcal{H}_q'(r_b) < 0 \qquad\Rightarrow\qquad \dot S(r_b) > 0, \tag{102}$$

or (from (35)) in terms of the local perturbations $\delta^{(\rho)}$, $\delta^{(\mathcal{H})}$:

$$\delta^{(\rho)}(r_b)\,\delta^{(\mathcal{H})}(r_b) < 0 \qquad\Rightarrow\qquad \dot S(r_b) > 0. \tag{103}$$

Either one of (102) or (103) allows us to test (at least qualitatively) the fulfillment of $\dot S > 0$ in arbitrary domains of generic LTB models by means of the analytic and qualitative results derived in [56, 39, 40]. We summarize in the next section some preliminary results (further results and more detail will be provided in [71]).

Besides the balance law (95), it is useful to compute time concavity/convexity of S given by the sign of $\ddot S$, which follows readily by applying (22) to (97):

$$\frac{\ddot S}{\mathcal{V}_q} = \frac{9\gamma_0}{2}\left\langle \rho[\mathbf{D}(\mathcal{H})]^2 + \frac{2}{3}\mathcal{H}_q\mathbf{D}(\rho)\mathbf{D}(\mathcal{H}) + \frac{4\pi}{9}[\mathbf{D}(\rho)]^2\right\rangle_q$$

♯ The condition $\rho'(r)\,\mathcal{H}'(r) < 0$ is also a sufficient condition if it holds for the full integration range $0 < r < r_b$ in the integrals in (101). On the other hand, conditions (102) and (103) only need to be evaluated at r_b. Notice that for monotonic profiles $\rho'(r_b) < 0$ and $\rho'(r_b) > 0$ respectively imply $\rho_q'(r_b) < 0$ and $\rho_q'(r_b) > 0$, but the converses are false, see [40].



Figure 6. Time slices and the expanding and collapsing singularities. The figure displays time slices t_1, t_2, t_3, t_4 (long dashed horizontal lines) in the (t, r) plane and the worldline of a comoving layer r = constant (short dashed vertical line) evolving from the a non–simultaneous big bang (thick lower curve $t_{\rm bb}(r)$) towards the collapsing singularity (thick upper curve $t_{\rm coll}(r)$) in an elliptic model. For t_1, t_2 we have necessarily $\mathcal{H}_q' < 0$ and $\rho_q' < 0$, whereas we have $\mathcal{H}_q' > 0$ and $\rho_q' < 0$ for t_1, t_2. Therefore, conditions (102) and (103) hold for slices near $t_{\rm coll}$ but fails for slices near $t_{\rm bb}$. The latter failure also occurs in hyperbolic models with non–simultaneous big bang that lack a collapsing singularity.

$$= \frac{3\gamma_0}{4}\left\langle \rho\,\sigma_{ab}\sigma^{ab} + \frac{1}{2\pi}\mathcal{H}_q\sigma_{ab}E^{ab} + \frac{1}{4\pi}E_{ab}E^{ab}\right\rangle_q, \qquad (104)$$

where we used (8a)–(8b), (44), (57b)–(57b) and (58a)–(58b) to eliminate the derivatives $\dot\rho$, $\dot\rho_q$, $\dot{\mathcal{H}}$, $\dot{\mathcal{H}}_q$ and Σ in terms of fluctuations $\mathbf{D}(\rho)$, $\mathbf{D}(\mathcal{H})$.

14. Conclusion and summary.

We have considered the q–average (the scalar average defined by (13)) in spherical comoving domains $\mathcal{D}[r]$ for the study of generic LTB models and their properties. This average generalizes the standard proper volume average of Buchert's formalism by the presence of a non–trivial weight factor, $\mathcal{F}$, that is suitable for these models (Buchert's average corresponds to the case $\mathcal{F} = 1$). By applying the q–average to LTB covariant scalars A, we obtained new dynamical variables (q–scalars) that can be defined as functionals $\langle A\rangle_q[r_b]$ (the average is assigned to a whole fixed but arbitrary domain) or as functions A_q (q–functions that depend on the average for domains with varying boundaries). The q–scalars are covariant, as they are expressible in terms of curvature invariants (see section 6), and also have very special and useful properties. Specifically, the q–scalars that are common with FLRW models (A_q and $\langle A\rangle_q[r_b]$ for $A = \rho$, $\mathcal{H}$, $\mathcal{K}$, Ω) identically satisfy FLRW fluid flow dynamics and their analytic forms mimic FLRW expressions that are commonly used in LTB void models that probe the possibility of explaining observations without resorting to dark energy (section 4.3). Also, the q–scalars determine (via Darmois matching conditions) a unique FLRW background state (section 4.4) that can be either a fictitious reference FLRW model that is different for each domain, or an actual background FLRW spacetime if a domain $\mathcal{D}[r_b]$ is smoothly matched to a FLRW region. Fluctuations and perturbations of the q–scalars can be defined by comparing them with non–averaged (*i.e.* local) covariant scalars. These fluctuations and perturbations are local ($\mathbf{D}(A)$, $\delta^{(A)}$) or non–local ($\mathbf{D}_{\rm NL}(A)$, $\delta_{\rm NL}^{(A)}$), depending on whether they are, respectively, defined for q–scalars that are functions or functionals. The following results are worth highlighting:

- All proper tensors (curvature and kinematic) characteristic of LTB models are expressible by linear combinations of g_{ab}, u^b and the common eigenframe of the shear and electric Weyl tensors ($\mathbf{e}_b^a$), with coefficients given by the q–scalars $A = \rho_q$, $\mathcal{H}_q$, $\mathcal{K}_q$, Ω_q and their relative local fluctuations (local perturbations) $\delta^{(\rho)}$, $\delta^{(\mathcal{H})}$, $\delta^{(\mathcal{K})}$, $\delta^{(\Omega)}$. In particular, all curvature tensors are expressible in terms of these linear combinations with coefficients given by the Ricci scalar $\mathcal{R}$ and its local fluctuations $\mathbf{D}(\mathcal{R})$ (section 6).
- Quadratic local fluctuations of ρ and $\mathcal{H}$ are equal to invariants that follow from contractions of the tensors involved in the dynamics of LTB models: the Ricci, electric Weyl and shear tensors. The q–averages of these fluctuations (local and non–local) are equal to statistical moments (variance, correlation and standard deviation) of ρ and $\mathcal{H}$ (see section 6).
- Local perturbations $\delta^{(A)}$ provide an elegant and invariant characterization of inhomogeneity in terms of the ratio of Weyl to Ricci curvature: $\Psi_2/\mathcal{R}$, where Ψ_2 is the only nonzero Newman–Penrose conformal invariant and $\mathcal{R}$ is the Ricci scalar, whereas relative Hubble scalar fluctuations $\delta^{(\mathcal{H})}$ do so in terms of the ratio of anisotropic vs. isotropic expansion: $\Sigma/\mathcal{H}$, where Σ is the eigenvalue of the shear tensor. All q–scalars can be given in terms of these scalar invariants (section 6).
- The evolution equations for the q–scalars and their perturbations (local and non–local) completely determine the dynamics of the models, and thus provide an alternative to the use of analytic solutions in the study and applications of the models. Although the resulting systems involve PDE's, they contain only time derivatives and the constraints are purely algebraic in nature and preserved in time. Hence, these equations can be treated effectively as ODE's (section 7).
- The basic difference with the standard proper volume average of Buchert's formalism is the absence of back–reaction correlation terms in the evolution equations associated with q–averages and their perturbations. Also, the relation between the averages of quadratic and higher order fluctuations and averages of scalar invariants and statistical moments of ρ and $\mathcal{H}$ does not occur with Buchert's average. We derive exact expressions for the Buchert's average of quadratic scalar invariants (see section 8).
- Since the q–variables behave effectively as FLRW scalars and their perturbations convey for every domain the (exact) deviation from FLRW dynamics, we can rigorously re–interpret LTB dynamics as the dynamics of exact spherical dust perturbations on a FLRW background defined by the q–scalars. As mentioned before, this background state is an abstract reference background defined at every domain by the continuity of the q–scalars under Darmois matching conditions (31a)–(31c). It is an actual FLRW spacetime only if a smooth matching with a FLRW region is considered in the context of "Swiss cheese" models (section 9).
- Both local and non–local fluctuations and perturbations can be either confined in a given domain $\mathcal{D}[r_b]$ (with or without assuming a "Swiss cheese" configuration through a matching with a FLRW region, see figures 1, 2 and 3), or an asymptotic perturbation for the case when $\mathcal{D}[r_b]$ becomes the whole time slice ${}^3\mathcal{T}[t]$ in the

limit $r_b \to \infty$ (see figure 4). The choice between confined and asymptotic perturbations depends on the boundary conditions of the specific problem or application that we may work out with LTB models. Confined perturbations (local or non–local) without a matching with FLRW are practical because they are easily applicable to study the inhomogeneity pattern of any generic LTB configuration, but are more abstract because the FLRW background state is a fictitious reference spacetime that changes for each domain. Swiss cheese configurations are more intuitive because the background state is a single (and non–fictitious) FLRW spacetime. They are useful if we wish to describe various inhomogeneous regions, as in Swiss cheese void models used to fit observations, but as as illustrated by panels (b) and (d) of figures 1, 2 and 3, their intuitiveness is offset by the artificial need to impose continuity of the perturbations (which must vanish in a non–ficticiuos background at finite radius). Asymptotic perturbations are more natural, and also practical if the radial profile of the scalars rapidly converges to a given set of FLRW values.

- The local perturbations provide a measure of inhomogeneity through the local magnitude of the radial gradients of covariant scalars, while non–local perturbations do so through the familiar notion of the "contrast" between local values of these scalars and a reference FLRW value defined as a q–average for each fixed whole domain, or for the whole slice in the asymptotic limit. Even if their evolution equations are more complicated, non–local perturbations are more intuitive than local ones, as they allow us to associate over and under densities with (respectively) positive or negative density amplitudes, while local perturbations yield the opposite sign. We have illustrated these points through the notion of the "amplitude" of the perturbations (see sections 10 and 11 and figure 5).
- The definition of non–local perturbations $\delta^{(A)}_{\rm NL}$ by the maps (75) and (76) provide (through the q–average and its relation to kinematic and curvature invariants) a rigorous and coordinate independent interpretation for simple examples of perturbations that are based on the notion of a contrast with respect to a FLRW background. These simple contrast perturbations are frequently introduced as ansatzes in many text–book and articles, for example, in the context of simple Newtonian models of structure formation (the "top hat" or "spherical collapse model" [67]) but also in linear perturbations [70] and in astrophysical applications of LTB models (see examples in [3, 4, 10]).
- Both local and non–local perturbations yield in the linear limit the familiar dust perturbations of linear theory in the comoving (or isochronous) gauge (section 12).

As a quick application, we examined in section 13 the definition of gravitational entropy based on the Liebler–Kullback functional from Information Theory, which has been proposed by Buchert and coworkers [42, 57, 58, 59]. Considering this functional as given in terms of q–averages yields the same conjecture obtained in these references, namely: a relation should exist between positive entropy production ($\dot S > 0$) and a negative correlation between fluctuations of ρ and $\mathcal{H}$. However, testing this conjecture is easier if we use q–averages, since: $\dot S$ is proportional to the q–average of the scalar invariant $\sigma_{ab}E^{ab}$, and also $\langle\rho\rangle_q$ and $\langle\mathcal{H}\rangle_q$ are expressible in closed analytic forms and the evolution equations for their fluctuations and perturbations (sections 7 and 8) do not involve complicated back–reaction terms. Hence, we were able to obtain the sufficient conditions for $\dot{S} > 0$ in terms of closed analytic expressions, such as (102) and (103), that can be tested in arbitrary domains of generic LTB models (a task that will be accomplished in future work [71]). Meanwhile, as "food for thought" we list the following preliminary results:

Entropy production is positive. Conditions (102) and (103) hold in the following cases:

- Late time evolution of elliptic models (see (A.3)) in time slices near the collapsing singularity. This result can be justified by qualitative arguments (see figure 6).
- Asymptotic time evolution ($t \to \infty$ for all r) of hyperbolic models (see (A.2)). This result follows from the fact that $\delta^{(\mathcal{H})} \to \delta^{(\mathcal{K})}/2$ holds in this limit for these models (because $\Omega_q \to 0$), and thus from table 1 of [40] we have $\delta^{(\rho)}\delta^{(\mathcal{H})} < 0$ for all initial conditions complying with regularity.

Entropy production is negative. Conditions (102) and (103) are violated in the following cases:

- Parabolic models (spatially flat models $\mathcal{K} = 0$, $\Omega_q = 1$, see (A.1)). Since $\delta^{(\mathcal{H})} = \delta^{(\rho)}/2$ holds for all t, and regularity conditions require $-1 < \delta^{(\mathcal{H})} \leq 0$ (see [39, 40]), then $\delta^{(\rho)}\delta^{(\mathcal{H})} > 0$ holds for all t.
- Early time evolution of elliptic and hyperbolic models with a non–simultaneous big bang. This result can be justified by qualitative arguments (see figure 6).

These results clearly indicate an important connection between the failure to fulfill a positive entropy production and specific conditions in which the density decaying modes dominate over the growing modes (parabolic models and near a non–simultaneous big bang [2, 3, 4, 30, 33]). This is an important theoretical feature worth a comprehensive examination (further detail is provided in [71]).

As a comparison with our approach and preliminary results, a recent perturbative study [59] of the Liebler–Kullback functional in the context of Buchert's average yields $\dot{S}$ only in terms of the average (Buchert's average) of the invariant scalar $C_{abcd}C^{abcd}$ (which is $8E_{ab}E^{ab}$ in LTB models), hence it excludes the contribution to inhomogeneity from the shear tensor (a spatial gradient of the velocity in the conformal Newtonian gauge) that comes from the scalar invariant $\sigma_{ab}E^{ab}$. Buchert and coworkers have not yet applied the Leibler–Kullback functional to generic LTB models in a non–perturbative manner (only a very simple example of an LTB model was considered in [58], but this model exhibits shell crossings). Although they obtained conditions that are analogous to (98), their main interest has been to provide a theoretical connection between the growth of structure complexity associated with $\dot{S} > 0$ and the dynamical implications of the back–reaction terms that appear in Buchert's formalism. Since the back–reaction terms of Buchert's formalism vanish for q–averages, our results provide a different theoretical perspective, namely: the connection between $\dot{S} > 0$ in (97) and $\ddot{S}$ in (104) associated with the growth of structure complexity and the statistical moments that arise from the q–average of quadratic invariant scalars ($E_{ab}E^{ab}$, $\sigma_{ab}E^{ab}$, $\sigma_{ab}\sigma^{ab}$) that vanish for FLRW models.

It is important to remark that the notion of a gravitational entropy connected to inhomogeneity (marked by invariant scalars of the Weyl tensor) bears a close relation to the concept of the "arrow of time" originally suggested by Penrose [72], and further explored by a number of authors [73, 74, 75, 76, 77]. The Weyl tensor scalar that emerges in this literature is $C_{abcd}C^{abcd}$ (proportional to $E_{ab}E^{ab}$ in LTB models) and no attempt is made in these papers to connect it with averages and fluctuations. As a contrast, the Leibler–Kullback functional is based on an averaging formalism that yields a direct connection between the growth of structure (inhomogeneity) and its associated fluctuations, which in the case of the q–average, are directly related to invariant contractions of σ_{ab} and E^{ab}, as well as with fluctuations and statistical variance and covariance moments of both ρ and $\mathcal{H}$ (which is a fluid velocity). Evidently, we have only considered this functional with the q–average applied on LTB models, and thus, we cannot exclude the possibility that contradictory results may happen with Buchert's average or in other types of models (as for example with the "arrow of time", see [75]), but the work we have developed here may provide a guideline on how to explore the notion of the arrow of time and its relation with the Liebler–Kullback functional in more general models.

While the q–average formalism cannot be used to study and understand the dynamical implications of back–reaction, it has resulted very helpful for understanding the properties and evolution of LTB models [39, 40], including the computation in these models of back–reaction terms and verification of the existence of "effective" acceleration in the context of Buchert's formalism [56]. Moreover, it does provide through its fluctuations and perturbations interesting connections between averaging, perturbation theory, invariant scalars and statistical correlations of ρ and $\mathcal{H}$, which signals a valuable theoretical insight on how the averaging process should work in any generic solution of Einstein's equations (at least in LRS spacetimes whose dynamics is reducible to scalar modes). In particular, since most formal and theoretical results obtained for LTB models can be readily applied to Szekeres models [62], the extension of our results to these models is currently under elaboration. In fact, the dynamical effect of q–scalars in LTB models may be applicable to even more general spacetimes. Specifically, we propose identifying fluid flow scalars that comply with FLRW dynamics (hopefully related to some weighed average), and then expressing local 1+3 scalars as fluctuations of the new scalars. The following step would be to explore if the 1+3 evolution equations given in terms of the new set of variables has the structure of evolution equations and constraints in the context of a formalism of exact perturbations on a FLRW background. We feel that this proposal is worth considering in future research.

Appendix A. Analytic solutions.

The metric function R in (1) can be obtained from the following parametric solutions of $(2a)$ which define parabolic, hyperbolic and elliptic models:

$$\textbf{Parabolic:}\; E=0,\quad R=(2M)^{1/3}\,\eta^2,\qquad t-t_{\rm bb}=\frac{2}{3}\,\eta^3, \tag{A.1}$$

$$\textbf{Hyperbolic:}\; E\geq 0,\quad R=\frac{M}{E}\,(\cosh\eta-1)\,,\quad t-t_{\rm bb}=\frac{M}{E^{3/2}}\,(\sinh\eta-\eta)\,, \tag{A.2}$$

$$\textbf{Elliptic:}\; -1<E\leq 0,\quad R=\frac{M}{|E|}\,(1-\cos\eta)\,,\quad t-t_{\rm bb}=\frac{M}{|E|^{3/2}}\,(\eta-\sin\eta)\,, \tag{A.3}$$

where $t_{\rm bb}=t_{\rm bb}(r)$, which emerges as an "integration constant", is the "big bang time" because it marks the coordinate locus of the central expanding curvature singularity $R(t,r)=0$ for $r\neq$ constant. The classification of (A.1)–(A.3) follows from the sign of E and governs the kinematic evolution of dust layers through the existence of a zero of $\dot R$ in $(2a)$: parabolic and hyperbolic layers perpetually expand and elliptic layers expand/collapse. Since $E = E(r)$, the models can exhibit, either the same kinematic pattern for all r, or "mixed" patterns that define models with regions of different kinematic behavior as E changes sign in specific ranges of r (see examples in [11]).

The solutions (A.1)–(A.3) allow for a complete description of the dynamics of the models once we select the free functions M, E, $t_{\rm bb}$, though only two of these functions are strictly necessary because the metric (1) in invariant under re–scalings $r = r(\bar{r})$ and thus any one of these free functions can be eliminated by a suitable choice of the radial coordinate. In particular, the expressions (2*b*), (3), (4), (5) and (6) for the basic covariant scalars $\rho, \mathcal{H}$, $\mathcal{K}$, Σ, $\mathcal{E}$ in (7) involve terms such as R' and $\dot{R}'$, which can be computed by implicit derivation of (A.1)–(A.3).

Appendix B. Functions of q–scalars are q–scalars.

The relations (23) allow us to express local scalars A in terms of q–scalars A_q through the differential inverse of the integral definition (13):

$$A = \langle A\rangle_q + \frac{\langle A\rangle_q'}{\mathcal{V}_q'/\mathcal{V}_q} = A_q + \frac{A_q'}{\mathcal{V}_q'/\mathcal{V}_q}, \tag{B.1}$$

where $\mathcal{V}_q'/\mathcal{V}_q = 3R'/R$ and $\mathcal{V}_q$ is defined in (94). As a consequence, any scalar expressible as a function of q–scalars is itself a q–scalar: if A_q and B_q are two q–scalars and $U = U(A_q, B_q)$, then

$$U' = \frac{3R'}{R}\left[\frac{\partial U}{\partial A_q}(A - A_q) + \frac{\partial U}{\partial B_q}(B - B_q)\right], \tag{B.2}$$

so that we can identify $Z_q = U(A_q, B_q)$ as the q–scalar whose corresponding "local" scalar Z is given by

$$Z = Z_q + \frac{\partial U}{\partial A_q}(A - A_q) + \frac{\partial U}{\partial B_q}(B - B_q) = Z_q + \frac{Z_q'}{3R'/R}, \tag{B.3}$$

where it is important to remark that (in general) we have $Z \neq U(A, B)$. It is straightforward to show that Z and Z_q above satisfy the integral definition (13), which basically states that a function of q–averages $U(\langle A\rangle_q, \langle B\rangle_q)$ is itself the q–average $\langle Z\rangle_q$ of a scalar Z given by (B.3), which in general does not coincide with $U(A, B)$ nor its q–average $\langle U(A, B)\rangle_q$.

Appendix C. Proof of (41) and (42).

The proof of (41) for $A = \mathcal{H}$ is equivalent to the proof that the identity $\langle W_{\mathcal{H}}(r, r_0)\rangle_q[r_b] = 0$ holds for every domain $\mathcal{D}[r_b]$, where the scalar $W_{\mathcal{H}}$ is given by:

$$W_{\mathcal{H}}(r, r_b) = [\mathbf{D}_{\rm NL}(\mathcal{H})]^2 - [\mathbf{D}(\mathcal{H})]^2 = (\,\mathcal{H}(r) - \langle\mathcal{H}\rangle_q[r_b]\,)^2 - (\,\mathcal{H}(r) - \mathcal{H}_q(r)\,)^2\,. \tag{C.1}$$

We expand $W_{\mathcal{H}}(r, r_b)$ and apply (21), so that $\langle -2\mathcal{H}(r)\langle\mathcal{H}\rangle_q[r_b] + \langle\mathcal{H}\rangle_q^2[r_b]\rangle_q[r_b] = -\langle\mathcal{H}\rangle_q^2[r_b]$ holds, leading to

$$\langle W_{\mathcal{H}}(r, r_b)\rangle_q[r_b] = -\langle\mathcal{H}\rangle_q^2[r_b] + \frac{1}{\mathcal{V}_q(r_b)}\int_0^{r_b}(2\mathcal{H}\mathcal{H}_q - \mathcal{H}_q^2)\,\mathcal{V}_q'\,{\rm d}\bar{r}. \tag{C.2}$$

where $\mathcal{V}_q$ is defined by (94), so that $\mathcal{V}_q' = 4\pi R^2 R'$. Considering that (3) and (19) imply $3\mathcal{H} = \theta = \dot{\mathcal{V}}_q'/\mathcal{V}_q'$ and $3\mathcal{H}_q = \theta_q = \dot{\mathcal{V}}_q/\mathcal{V}_q$. Inserting these identities into (C.2) yields the desired result:

$$\langle W_{\mathcal{H}}(r,r_b)\rangle_q[r_b] \;=\; -\,\langle\mathcal{H}\rangle_q^2[r_b] + \frac{1}{\mathcal{V}_q(r_b)}\int_0^{r_b}\left(\frac{\dot{\mathcal{V}}_q^2}{9\mathcal{V}_q}\right)'\,\mathrm{d}\bar{r} = -\langle\mathcal{H}\rangle_q^2[r_b] + \mathcal{H}_q^2(r_b) = 0, \tag{C.3}$$

since $\langle\mathcal{H}\rangle_q[r_b]$ coincides with $\mathcal{H}_q$ at the domain boundary $r = r_b$.

For $A = \rho$ the proof of (41) is equivalent to proving $W_\rho(r,r_b)\rangle_q[r_b] = 0$, where

$$W_\rho(r,r_b) = [\mathbf{D}_{\rm NL}(\rho)]^2 - [\mathbf{D}(\rho)]^2 = (\,\rho(r) - \langle\rho\rangle_q[r_b]\,)^2 - (\,\rho(r) - \rho_q(r)\,)^2\,. \tag{C.4}$$

Expanding $W_\rho(r,r_b)$, applying (21) and substituting $\rho = M'/\mathcal{V}_q'$ and $\rho_q = M/\mathcal{V}_q$ from (2*b*) and (19), yields a relation similar to (C.2):

$$\begin{aligned}\langle W_\rho(r,r_b)\rangle_q[r_b] \;&=\; -\,\langle\rho\rangle_q^2[r_b] + \frac{1}{\mathcal{V}_q(r_b)}\int_0^{r_b}(2\rho\rho_q - \rho_q^2)\,\mathcal{V}_q'\,\mathrm{d}\bar{r},\\ &=\; -\,\langle\rho\rangle_q^2[r_b] + \frac{1}{\mathcal{V}_q(r_b)}\int_0^{r_b}\left(\frac{M^2}{\mathcal{V}_q}\right)'\,\mathrm{d}\bar{r} = -\langle\rho\rangle_q^2[r_b] + \rho_q^2(r_b) = 0.\end{aligned} \tag{C.5}$$

The proof of (42) for $A, B = \rho$, $\mathcal{H}$ follows along similar lines: it is equivalent to proving that: $\langle\, W_{\rho\mathcal{H}}(r,r_b)\,\rangle_q[r_b] = 0$ holds, where:

$$\begin{aligned}&W_{\rho\mathcal{H}}(r,r_b) = \mathbf{D}_{\rm NL}(\mathcal{H})\,\mathbf{D}_{\rm NL}(\rho) - \mathbf{D}(\mathcal{H})\,\mathbf{D}(\mathcal{H})\\ &= (\,\rho(r) - \langle\rho\rangle_q[r_b]\,)\,(\,\mathcal{H}(r) - \langle\mathcal{H}\rangle_q[r_b]\,) - (\,\rho(r) - \rho_q(r)\,)\,(\,\mathcal{H}(r) - \mathcal{H}_q(r)\,)\,.\end{aligned} \tag{C.6}$$

Substitution of the forms for $\mathcal{H}$, $\mathcal{H}$, ρ, ρ_q given before yields the desired result:

$$\begin{aligned}\langle W_{\rho\mathcal{H}}(r,r_b)\rangle_q[r_b] &= -\langle\rho\rangle_q[r_b]\langle\mathcal{H}\rangle_q[r_0] + \frac{1}{\mathcal{V}_q(r_b)}\int_0^{r_b}(\rho\mathcal{H}_q + \rho_q\mathcal{H} - \rho_q\mathcal{H}_q)\,\mathcal{V}_q'\,\mathrm{d}\bar{r}\\ &= -\langle\rho\rangle_q[r_b]\langle\mathcal{H}\rangle_q[r_b] + \frac{1}{\mathcal{V}_q(r_b)}\int_0^{r_b}\left(\frac{M\dot{\mathcal{V}}_q}{3\mathcal{V}_q}\right)'\,\mathrm{d}\bar{r}\\ &= -\langle\rho\rangle_q[r_b]\langle\mathcal{H}\rangle_q[r_b] + \rho_q(r_b)\mathcal{H}_q(r_b) = 0.\end{aligned} \tag{C.7}$$

Appendix D. Proper curvature and kinematic tensors.

We show in this appendix that all relevant (curvature and kinematic) proper tensors characteristic of LTB models are completely determined by 4 invariant scalars: the Ricci scalar, $\mathcal{R} = \mathcal{R}_c^c$, the only nonzero Newman–Penrose conformal invariant, Ψ_2, the Hubble expansion scalar $\mathcal{H} = \theta/3$ and the eigenvalue of the shear tensor, Σ.

The Riemann and Weyl tensors are given by the following expressions:

$$\mathcal{R}_{cd}^{ab} = C_{cd}^{ab} - \frac{1}{2}\mathcal{R}\delta_{[c}^{a}\delta_{d]}^{b} + 2\delta_{[c}^{[a}\mathcal{R}_{d]}^{b]}, \qquad C_{cd}^{ab} = \delta_{[c}^{[a}E_{d]}^{b]} + 2u_{[c}E_{d]}^{[a}u^{b]}, \tag{D.1}$$

where square bracketed indices indicate anti–simetrization. The Ricci and electric Weyl tensors, $\mathcal{R}_d^b$ and E_d^b follow from contraction of the tensors in (D.1):

$$\mathcal{R}_d^b = \mathcal{R}_{cd}^{bc} = \frac{1}{2}\mathcal{R}\left(h_d^b + u^b u_d\right), \qquad E_d^b = u_a u_c C_d^{abc} = \Psi_2\,\mathbf{e}_d^b, \tag{D.2}$$

where $\mathbf{e}^{b}_{d} = \delta^{b}_{d} - 3n^{b}n_{d}$ is the common eigenframe for E^{b}_{d} and the shear tensor σ^{b}_{d} defined in (5) and (6). By combining (D.1) and (D.2) we get

$$\mathcal{R}^{ab}_{cd} = C^{ab}_{cd} + \frac{1}{3}\mathcal{R}\left(3\delta^{[a}_{[c}\delta^{b]}_{d]} + 6\delta^{[a}_{[c}u^{b]}u_{d]} - \delta^{a}_{[c}\delta^{b}_{d]}\right), \tag{D.3}$$

$$C^{ab}_{cd} = \Psi_2\left(\delta^{[a}_{[c}\mathbf{e}^{b]}_{d]} + 2u_{[c}\mathbf{e}^{[a}_{d]}u^{b]}\right) = \Psi_2\left(h^{[a}_{[c} - 3u_{[c}u^{[a}\right)\mathbf{e}^{b]}_{d]}, \tag{D.4}$$

Hence, all curvature proper tensors are determined by two invariant scalars: $\mathcal{R}$ and Ψ_2, which relate to the density and its local fluctuation by $\mathcal{R} = 8\pi\rho$ and $\Psi_2 = -\mathcal{E} = (4\pi/3)(\rho - \rho_q) = (4\pi/3)\mathbf{D}(\rho)$.

The kinematic proper tensors are the shear and "expansion" tensors: σ_{ab}, $\mathcal{H}_{ab}$. Both are expressible in terms of the invariant scalars $\mathcal{H} = \theta/3 = \tilde{\nabla}_{a}u^{a}/3$ and $\Sigma = \sigma_{ab}\mathbf{e}^{ab}$:

$$\sigma_{ab} = \Sigma\,\mathbf{e}_{ab}, \qquad \mathcal{H}_{ab} = \mathcal{H}h_{ab} + \Sigma\,\mathbf{e}_{ab}, \tag{D.5}$$

where the explicit forms of $\mathcal{H}$ and Σ are given in (3) and (5). Notice that $\Sigma = -\mathbf{D}(\mathcal{H})$ is a local fluctuation of $\mathcal{H}$.

References

[1] Lemaître G 1933 *Ann. Soc. Sci. Brux.* A **53** 51. See reprint in Lemaître G 1997 *Gen. Rel. Grav.* **29** 5; Tolman R C 1934 *Proc. Natl Acad. Sci.* **20** 169; Bondi H 1947 *Mon. Not. R. Astron. Soc.* **107** 410.

[2] Krasiński A, *Inhomogeneous Cosmological Models*, Cambridge University Press, 1998.

[3] Plebanski J and Krasinski A, *An Introduction to General Relativity and Cosmology*, Cambridge University Press, 2006.

[4] K. Bolejko, A. Krasiński, C. Hellaby, M.-N. Célérier, Structures in the Universe by exact methods: formation, evolution, interactions, Cambridge University Press, Cambridge 2009

[5] Celerièr M N 2007 *New Advances in Physics* **1** 29 (*Preprint* `arXiv:astro-ph/0702416`)

[6] Krasiński A and Hellaby C 2002 *Phys Rev* D **65** 023501

[7] Krasiński A and Hellaby C 2004 *Phys Rev* D **69** 023502

[8] Krasiński A and Hellaby C 2004 *Phys Rev* D **69** 043502

[9] Hellaby C and Krasiński A 2006 *Phys Rev* D **73** 023518

[10] Bolejko K Krasiński A and Hellaby C 2005 *MNRAS* **362** 213 (*Preprint* `arXiv:gr-qc/0411126`)

[11] Matravers D R and Humphreys N P 2001 *Gen. Rel. Grav.* **33** 53152; Humphreys N P, Maartens R and Matravers D R 1998 Regular spherical dust spacetimes (*Preprint* `gr-qc/9804023v1`)

[12] Bolejko K Krasinski A and Hellaby C 2005 *Mon Not Roy Astron Soc* **362** 213-228 (*Preprint* `arXiv:0411126v2[gr-qc]`)

[13] Bolejko K Celerier M N and Krasinski A 2011 *Class. Quant. Grav.* **28** 164002;

[14] Pascual–Sánchez J F 1999 *Mod. Phys. Lett.* A **14** 1539; Sugiura N K and Harada T 1999 *Phys Rev* D 60 103508; Celerièr M N 2000 *Astron. Astrophys.* **353** 63; Tomita K 2001 *MNRAS* **326** 287; Iguchi H, Nakamura T and Nakao K 2002 *Prog. Theor. Phys.* **108** 809; Schwarz D J 2002 Accelerated expansion without dark energy (*Preprint* `arXiv:astro-ph/0209584v2`);

[15] Apostolopoulos P *et al* 2006 *JCAP* **P06** 009; Kai T, Kozaki H, Nakao K, Nambu Y and Yoo C M 2007 *Prog. Theor. Phys.* **117** 229-240 (*Preprint* `arXiv:gr-qc/0605120`); Mattsson T and Ronkainen M 2008 *JCAP* **0802** 004 (*Preprint* `arXiv:astro-ph/0708.3673v2`); Bolejko K and Andersson L 2008 *JCAP* **10** 003 (*Preprint* `arXiv:0807.3577`); Rasanen S 2006 *Class. Quant. Grav.* **23** 1823-1835;

[16] Moffat J W 2006 *J. Cosmol. Astropart. Phys.* JCAP(2006)001;

[17] Kolb E W, Matarrese S, Notari A and Riotto A 2005 *Phys Rev* D **71** 023524 (*Preprint* `arXiv:hep-ph/0409038v2`); Marra V, Kolb E W and Matarrese S 2008 *Phys Rev* D **77** 023003; Marra V, Kolb E W, Matarrese S and Riotto A 2007 *Phys Rev* D **76** 123004.

[18] García–Bellido J and Troels H 2008 **JCAP** 0804:003 (*Preprint* `gr-qc/0802.1523v3 [astro-ph]`)

[19] Alnes H, Amazguioui M and Gron O 2006 *Phys Rev* D **73** 083519; Alnes H and Amazguioui M 2006 *Phys Rev* D **74** 103520; Alnes H and Amazguioui M 2006 *Phys Rev* D **75** 023506

[20] Enqvist K and Mattsson T 2007 *JCAP* **0702** 019 (*Preprint* `arXiv:astro-ph/0609120v4`); Enqvist K 2008 *Gen. Rel. Grav.* **40** 451-466 (*Preprint* `arXiv:0709.2044`)

[21] Biswas T Notari A and Valkenburg W 2010 *JCAP* **11** 030 (*Preprint* `arXiv:1007.3065`)

[22] February S *et al* 2010 *MNRAS* **405** 2231–2242 (*Preprint* arXiv:0909.1479v2[astro-ph CO])

[23] Marra V and Notari A 2011 *Class. Quant. Grav.* **28** 164004 (*Preprint* arXiv:1102.1015)

[24] Eardley D M 1974 *Commun Math Phys* **37** 287; Eardley D M and Smarr L 1979 *Phys Rev* D **19** 2239; Dyer C C 1979 *MNRAS* **189** 189; Waugh B and Lake K 1988 *Phys Rev* D **38** 1315; Waugh B and Lake K 1989 *Phys Rev* D **40** 2137; Lemos J P S 1991 *Phys Lett* A **158** 279

[25] Joshi P S and Dwivedi I H 1993 *Phys Rev* D **47** 5357; Joshi P S and Singh T P 1995 *Phys Rev* D **51** 6778; Dwivedi I H and Joshi P S 1997 *Class. Quant. Grav.* **47** 5357

[26] Vaz C, Witten L and Singh T P 2001 *Phys Rev* D **63** 104020; Kiefer C, Mueller-Hill, Vaz C 2006 *Phys Rev* D **73** 044025; Bojowald M, Harada T and Tibrewala R 2008 *Phys Rev* D **78** 064057

[27] Ellis G F R and Bruni M 1989 *Phys Rev* D **40** 1804

[28] Bruni M, Dunsby P K S and Ellis G F R 1992 *Astroph. J.* **395** 34–53

[29] Ellis G F R and van Elst H 1998 Cosmological Models (Cargèse Lectures 1998) *Preprint* arXiv gr-qc/9812046 v4

[30] Zibin J P 2008 *Phys Rev* D**78** 043504 [arXiv:0804.1787]

[31] Dunsby P *et al* (*Preprint* arXiv:1002.2397v1 [astro-ph CO])

[32] van Elst H and Ellis G F R 1996 *Class Quantum Grav* **13** 1099-1128 (*Preprint* arXiv:gr-qc/9510044)

[33] Wainwright J and Andrews S 2009 *Class.Quant.Grav.*,**26**, 085017

[34] Sussman R A and García–Trujillo L 2002 *Class.Quant.Grav.* **19** 2897-2925.

[35] Sussman R A Quasi-local variables and inhomogeneous cosmological sources with spherical symmetry 2008 *AIP Conf.Proc.* **1083** 228-235 Preprint arXiv:0810.1120.

[36] Sussman R A 2009 *Phys Rev* D **79** 025009 (*Preprint* arXiv:arXiv:0801.3324 [gr-qc])

[37] Sussman R A 2008 *Class Quantum Grav.* **25** 015012 Preprint arXiv:grqc/0709.1005

[38] Sussman R A and Izquierdo G 2011 *Class Quantum Grav.* **28** 045006 Preprint arXiv:grqc/1004.0773

[39] Sussman R A 2010 *Gen Rel Grav* **42** 2813–2864 (*Preprint* arXiv:1002.0173 [gr-qc])

[40] Sussman R A 2010 *Class.Quant.Grav.* **27** 175001 (*Preprint* arXiv:1005.0717 [gr-qc])

[41] Buchert T, 2000 *Gen. Rel. Grav.* **32** 105; Buchert T, 2000 *Gen.Rel.Grav.* **32** 306-321; Buchert T 2001 *Gen. Rel. Grav.* **33** 1381-1405; Ellis G F R and Buchert T 2005 *Phys.Lett.* A**347** 38-46; Buchert T and Carfora M 2002 *Class.Quant.Grav.* **19** 6109-6145; Buchert T 2006 *Class. Quantum Grav.* 23 819; Buchert T, Larena J and Alimi J M 2006 *Class. Quantum Grav.* 23 6379; Buchert T 2005 *Class. Quantum Grav.* **22** L113–L119; Buchert T 2006 *Class. Quantum Grav.* **23** 817–844 (*Preprint* arXiv:gr-qc/0509124)

[42] Buchert T 2008 *Gen. Rel. Grav.* **40** 467–527 (*Preprint* arXiv:xxxx.xxxx)

[43] Buchert T 2011 *Class. Quantum Grav.* **28** 164007 (*Preprint* arXiv:1103.2016)

[44] Zalaletdinov R M, *Averaging Problem in Cosmology and Macroscopic Gravity*, Online Proceedings of the Atlantic Regional Meeting on General Relativity and Gravitation, Fredericton, NB, Canada, May 2006, ed. R.J. McKellar (*Preprint* arXiv:gr-qc/0701116)

[45] Coley A A and Pelavas N 2007 Phys.Rev. D **75** 043506; Coley A A, Pelavas N and Zalaletdinov R M 2005 *Phys.Rev.Lett.* **95** 151102

[46] Buchert T and Carfora M 2002 *Class. Quantum Grav.* **19** (*Preprint* arXiv:gr-qc/0210037); Paranjape A and Singh T P 2008 *Gen.Rel.Grav.* **40** 139-157 (*Preprint* arXiv:astro-ph/0609481v4); Paranjape A and Singh T P 2007 *Phys.Rev.* D **76** 044006 (*Preprint* arXiv:gr-qc/0703106v3); Paranjape A 2008 *Int.J.Mod.Phys.* D **17** 597-601 (*Preprint* arXiv:0705.2380v1 [gr-qc]); Marozzi G 2011 *JCAP* **01** 012 (*Preprint* arXiv:1011.4921 [gr-qc])

[47] Buchert T 2011 *Classical and Quantum Gravity* **28** 164007; Clarkson C Ellis G F R Larena J and Umeh O 2011 *Reports on Progress in Physics*, (*Preprint* arXiv:1109.2314); Clarkson C Ananda K and Larena J 2009 *Phys Rev* D **80** 083525 (*Preprint* arXiv:0907.3377); Rasänen S 2008 *JCAP* **0804** 027 (*Preprint* arXiv:0801.2692); Paranjape A 2008 *Phys.Rev.D* **78** 063522 (*Preprint* arXiv:0806.2755v2 [astro-ph]); Paranjape A and Singh T P 2008 *Phys.Rev.Lett.* **101** 181101 (*Preprint* arXiv:0806.3497v3 [astro-ph]); Mattsson T and Ronkainen M 2008 *JCAP* **02** 004; Buchert T 2006 *Astron Astrophys* **454** 415-422 (*Preprint* arXiv:astro-ph/0601513)

[48] Kolb E *Classical and Quantum Gravity* **28** 164009; Clarkson C and Umeh O 2011 *Classical and Quantum Gravity* **xx** (*Preprint* arXiv:1105.1886) Larena J Alimi J M Buchert T Kunz M and Corasaniti P S 2009 *Phys.Rev.* D **79** 083011 (*Preprint* arXiv:0808.1161); Kolb E W Matarrese S and Riotto A 2006 *NewJ.Phys.* **8** 322 (*Preprint* arXiv:astro-ph/0506534);

[49] Buchert T 2005 *Class Quant Grav* **22** L113-L119 (*Preprint* arXiv:gr-qc/0507028); Ellis GFR and Buchert T 2005 *Phys Lett* A **347** 38-46 (*Preprint* arXiv:gr-qc/0506106); Räsänen S 2011 *Class Quantum Grav* **28** 164008 (*Preprint* arXiv:1102.0408)

[50] Buchert T and Obadia N 2011 *Class Quant Grav* **28** 162002 (*Preprint* arXiv:1010.4512); Wiegand A and Buchert T 2010 *Phys Rev* D**82** 023523 (*Preprint* arXiv:1002.3912); Roy X and Buchert T 2010 *Class Quant Grav* **27** 175013 (*Preprint* arXiv:0909.4155)

[51] Chuang C H, Gu J A and Hwang W Y P 2005 *Class.Quant.Grav.*,**25**, 175001 Preprint astro-ph/0512651

[52] Paranjape A and Singh T P 2006 *Class Quant Grav* **23**, 69556969

[53] Mattsson M and Mattsson T 2010 *JCAP* **10** 021; Mattsson M and Mattsson T 2011 *JCAP* **05** 003;

[54] Sussman R A 2008 On spatial volume averaging in Lematre–Tolman–Bondi dust models. Part I: back reaction, spacial curvature and binding energy (*Preprint* arXiv:0807.1145 [gr-qc])

[55] Sussman R A 2010 *AIP Conf Proc* **1241** 1146-1155 (*Preprint* arXiv:0912.4074 [gr-qc])

[56] Sussman R A 2011 *Class Quantum Grav* **28** 235002 (*Preprint* arXiv:1102.2663v1 [gr-qc])

[57] Hosoya A Buchert T and Morita M 2004 *Phys Rev Lett* **92**, 141302-14 (*Preprint* arXiv:gr-qc/0402076)

[58] Morita M Buchert T Hosoya A and Li N 2010 *AIP Conf.Proc.* **1241** 1074-1082 (*Preprint* arXiv:1011.5604)

[59] Li N Buchert T Hosoya A Morita M and Schwarz D J 2012 "Relative information entropy and Weyl curvature of the inhomogeneous Universe" (*Preprint* arXiv:1208.3376) arXiv:1208.3376

[60] Hayward S A 1996 *Phys Rev* D **53** 1938 (*Preprint* ArXiv gr-qc/9408002); Hayward S A 1998 *Class Quantum Grav* **15** 31473162 (*Preprint* ArXiv gr-qc/9710089v2)

[61] Misner C and Sharp D H 1964 *Phys Rev* B **136** 571

[62] Sussman R A and Bolejko K 2012 *Class Quant Grav* **29** 065018 (*Preprint* ArXiv 1109.1178)

[63] Liddle A R and Lyth D H 1993 *Phys Rept* **231** 1-105 (*Preprint* arXiv:astro-ph/9303019v1). See sections 2.3 and 2.4

[64] Malik K A and Wands D 2009 *Phys Rept* **475** 1-51.

[65] Stewart J M and Walker M 1974 *Proc. R. Soc. London* A **341** 49

[66] Bardeen J 1980 *Phys Rev* D **22** 1882; Bardeen J, Steinhardt P and Turner M S 1983 *Phys Rev* D **28** 679

[67] Padmanabhan T 2002 *Theoretical Astrophysics, Volume III: Galaxies and Cosmology* (Cambridge University Press). See equation 5.22, page 278

[68] Kasai M 1992 *Phys Rev Lett* **69** 2330; Kasai M 1993 *Phys Rev* D **47** 3214

[69] Ishak M and Peel A 2012 *Phys Rev* D **85** 083502 (*Preprint* arXiv:1104.2590)

[70] Hwang J and Noh H 2006 *Phys Rev* D**73** 044021 (*Preprint* arXiv:astro-ph/0601041v1)

[71] Sussman R A 2012 "Gravitational entropy in LTB and Szekeres models" (in preparation).

[72] Penrose R 1979 in *General Relativity, an Einstein Centenary Survey.* Edited by Hawking S W and Israel W, Cambridge University Press.

[73] Wainwright J 1984 *Gen Rel Grav* **16** 657

[74] Bonnor W B 1986 *Class Quantum Grav* **3** 495

[75] Bonnor W B 1987 *Phys Lett* A **122** 305

[76] Gron O and Hervik S 2002 "The Weyl curvature conjecture" (*Preprint* gr-qc/0205026v1)

[77] Pelavas N and Lake K 2000 *Phys Rev* D **62** 044009